

## Product datasheet for KN208818LP

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### **BCDIN3D Human Gene Knockout Kit (CRISPR)**

#### **Product data:**

**Product Type:** Knockout Kits (CRISPR)

**Format:** 2 gRNA vectors, 1 Luciferase-Puro donor, 1 scramble control

**Donor DNA:** Luciferase-Puro

Symbol: BCDIN3D Locus ID: 144233

**Components: KN208818G1**, BCDIN3D gRNA vector 1 in pCas-Guide CRISPR vector (GE100002)

KN208818G2, BCDIN3D gRNA vector 2 in pCas-Guide CRISPR vector (GE100002)

KN208818LPD, donor DNA containing left and right homologous arms and Luciferase-Puro

functional cassette.

**GE100003**, scramble sequence in pCas-Guide vector

**Disclaimer:** These products are manufactured and supplied by OriGene under license from ERS. The kit is

designed based on the best knowledge of CRISPR technology. The system has been functionally validated for knocking-in the cassette downstream the native promoter. The efficiency of the knock-out varies due to the nature of the biology and the complexity of the

experimental process.

 RefSeq:
 NM 181708

 UniProt ID:
 Q7Z5W3

Synonyms: BCDIN3 domain containing

**Summary:** This gene encodes an RNA methyltransferase which belongs to the rossmann fold

methyltransferase family, and serves as a 5'-methylphosphate capping enzyme that is

specific for cytoplasmic histidyl tRNA. The encoded protein contains an Sadenosylmethionine binding domain and uses the methyl group donor, Sadenosylmethionine binding domain and uses the methyl group donor, Sadenosylmethionine binding domain and uses the methyl group donor, Sadenosylmethionine binding domain and uses the methyl group donor, Sadenosylmethionine binding domain and uses the methyl group donor, Sadenosylmethionine binding domain and uses the methyl group donor, Sadenosylmethionine binding domain and uses the methyl group donor, Sadenosylmethionine binding domain and uses the methyl group donor, Sadenosylmethionine binding domain and uses the methyl group donor, Sadenosylmethionine binding domain and uses the methyl group donor, Sadenosylmethionine binding domain and uses the methyl group donor, Sadenosylmethionine binding domain and uses the methyl group donor, Sadenosylmethionine binding domain and uses the methyl group donor, Sadenosylmethionine binding domain and uses the methyl group donor, Sadenosylmethionine binding domain and uses the methyl group donor, Sadenosylmethionine binding domain and uses the methyl group donor, Sadenosylmethionine binding domain and uses the methyl group donor, Sadenosylmethionine binding domain and uses the methyl group donor and use the methyl group donor and use

adenosylmethionine. This gene is overexpressed in breast cancer cells, and is related to the tumorigenic phenotype and poor prognosis of breast cancer. The encoded protein is thought to promote the cellular invasion of breast cancer cells, by downregulating the expression of tumor suppressor miRNAs through the dimethylation of the 5-monophosphate of the

PNA For State A 20471

corresponding precursor miRNAs. [provided by RefSeq, Apr 2017]





# **Product images:**

#### Donor Vector Edited Chromosome



RFP, Luc, and mBFP will be under native gene promoter