

# Product datasheet for KN216946RB

# alpha A Crystallin (CRYAA) Human Gene Knockout Kit (CRISPR)

## **Product data:**

Product Type:	Knockout Kits (CRISPR)
Format:	2 gRNA vectors, 1 RFP-BSD donor, 1 scramble control
Donor DNA:	RFP-BSD
Symbol:	alpha A Crystallin
Locus ID:	1409
Components:	<ul> <li>KN216946G1, alpha A Crystallin gRNA vector 1 in pCas-Guide CRISPR vector (GE100002)</li> <li>KN216946G2, alpha A Crystallin gRNA vector 2 in pCas-Guide CRISPR vector (GE100002)</li> <li>KN216946RBD, donor DNA containing left and right homologous arms and RFP-BSD functional cassette.</li> <li>GE100003, scramble sequence in pCas-Guide vector</li> </ul>
Disclaimer:	These products are manufactured and supplied by OriGene under license from ERS. The kit is designed based on the best knowledge of CRISPR technology. The system has been functionally validated for knocking-in the cassette downstream the native promoter. The efficiency of the knock-out varies due to the nature of the biology and the complexity of the experimental process.
RefSeq:	<u>NM 000394</u> , <u>NM 001363766</u>
UniProt ID:	<u>P02489</u>
Synonyms:	CRYA1; CTRCT9; HSPB4

View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### STANDARIA STANDARD CRYAA) Human Gene Knockout Kit (CRISPR) – KN216946RB 🖉

Summary: Mammalian lens crystallins are divided into alpha, beta, and gamma families. Alpha crystallins are composed of two gene products: alpha-A and alpha-B, for acidic and basic, respectively. Alpha crystallins can be induced by heat shock and are members of the small heat shock protein (HSP20) family. They act as molecular chaperones although they do not renature proteins and release them in the fashion of a true chaperone; instead they hold them in large soluble aggregates. Post-translational modifications decrease the ability to chaperone. These heterogeneous aggregates consist of 30-40 subunits; the alpha-A and alpha-B subunits have a 3:1 ratio, respectively. Two additional functions of alpha crystallins are an autokinase activity and participation in the intracellular architecture. The encoded protein has been identified as a moonlighting protein based on its ability to perform mechanistically distinct functions. Alpha-A and alpha-B gene products are differentially expressed; alpha-A is preferentially restricted to the lens and alpha-B is expressed widely in many tissues and organs. Defects in this gene cause autosomal dominant congenital cataract (ADCC). [provided by RefSeq, Jan 2014]

### **Product images:**



RFP, Luc, and mBFP will be under native gene promoter

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US