

Product datasheet for KN304781

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Rockville, MD 20850, US

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Dpp4 Mouse Gene Knockout Kit (CRISPR)

Product data:

Product Type: Knockout Kits (CRISPR)

Format: 2 gRNA vectors, 1 GFP-puro donor, 1 scramble control

Donor DNA: GFP-puro

Dpp4 Symbol: Locus ID: 13482

KN304781G1, Dpp4 gRNA vector 1 in pCas-Guide CRISPR vector (GE100002), Target Sequence: Components:

GGTTGCCTCCAGGAAAACTG

KN304781G2, Dpp4 gRNA vector 2 in pCas-Guide CRISPR vector (GE100002), Target Sequence:

CCCGACCCTCTGGTCCAGGG

KN304781D, donor DNA containing left and right homologous arms and GFP-puro functional

cassette.

GE100003, scramble sequence in pCas-Guide vector

RefSeq: NM 001159543, NM 010074

UniProt ID: P28843

Synonyms: Cd26; Dpp-4; THAM

Summary: Cell surface glycoprotein receptor involved in the costimulatory signal essential for T-cell

> receptor (TCR)-mediated T-cell activation. Acts as a positive regulator of T-cell coactivation, by binding at least ADA, CAV1, IGF2R, and PTPRC. Its binding to CAV1 and CARD11 induces T-cell proliferation and NF-kappa-B activation in a T-cell receptor/CD3-dependent manner. Its interaction with ADA also regulates lymphocyte-epithelial cell adhesion. In association with FAP is involved in the pericellular proteolysis of the extracellular matrix (ECM), the migration and invasion of endothelial cells into the ECM. May be involved in the promotion of lymphatic endothelial cells adhesion, migration and tube formation. When overexpressed, enhanced cell proliferation, a process inhibited by GPC3. Acts also as a serine exopeptidase with a dipeptidyl peptidase activity that regulates various physiological processes by cleaving peptides in the circulation, including many chemokines, mitogenic growth factors, neuropeptides and peptide hormones. Removes N-terminal dipeptides sequentially from

polypeptides having unsubstituted N-termini provided that the penultimate residue is

proline.[UniProtKB/Swiss-Prot Function]





Product images:

