

Product datasheet for **KN307617LP**

Hdac2 Mouse Gene Knockout Kit (CRISPR)

Product data:

Product Type:	Knockout Kits (CRISPR)
Format:	2 gRNA vectors, 1 Luciferase-Puro donor, 1 scramble control
Donor DNA:	Luciferase-Puro
Symbol:	Hdac2
Locus ID:	15182
Components:	KN307617G1 , Hdac2 gRNA vector 1 in pCas-Guide CRISPR vector (GE100002) KN307617G2 , Hdac2 gRNA vector 2 in pCas-Guide CRISPR vector (GE100002) KN307617LPD , donor DNA containing left and right homologous arms and Luciferase-Puro functional cassette. GE100003 , scramble sequence in pCas-Guide vector
Disclaimer:	These products are manufactured and supplied by OriGene under license from ERS. The kit is designed based on the best knowledge of CRISPR technology. The system has been functionally validated for knocking-in the cassette downstream the native promoter. The efficiency of the knock-out varies due to the nature of the biology and the complexity of the experimental process.
RefSeq:	NM_008229
UniProt ID:	P70288
Synonyms:	D10Wsu179e; mRPD3; YAF1; Yy1bp



[View online »](#)

Summary:

Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes (By similarity). Forms transcriptional repressor complexes by associating with MAD, SIN3, YY1 and N-COR. Interacts in the late S-phase of DNA-replication with DNMT1 in the other transcriptional repressor complex composed of DNMT1, DMAP1, PCNA, CAF1. Deacetylates TSHZ3 and regulates its transcriptional repressor activity. Component of a RCOR/GFI/KDM1A/HDAC complex that suppresses, via histone deacetylase (HDAC) recruitment, a number of genes implicated in multilineage blood cell development. May be involved in the transcriptional repression of circadian target genes, such as PER1, mediated by CRY1 through histone deacetylation. Involved in MTA1-mediated transcriptional corepression of TFF1 and CDKN1A.[UniProtKB/Swiss-Prot Function]

Product images:
