

Product datasheet for **KN317625BN**

Tlr2 Mouse Gene Knockout Kit (CRISPR)

Product data:

Product Type:	Knockout Kits (CRISPR)
Format:	2 gRNA vectors, 1 mBFP-Neo donor, 1 scramble control
Donor DNA:	mBFP-Neo
Symbol:	Tlr2
Locus ID:	24088
Components:	KN317625G1 , Tlr2 gRNA vector 1 in pCas-Guide CRISPR vector (GE100002) KN317625G2 , Tlr2 gRNA vector 2 in pCas-Guide CRISPR vector (GE100002) KN317625BND , donor DNA containing left and right homologous arms and mBFP-Neo functional cassette. GE100003 , scramble sequence in pCas-Guide vector
Disclaimer:	These products are manufactured and supplied by OriGene under license from ERS. The kit is designed based on the best knowledge of CRISPR technology. The system has been functionally validated for knocking-in the cassette downstream the native promoter. The efficiency of the knock-out varies due to the nature of the biology and the complexity of the experimental process.
RefSeq:	NM_011905
UniProt ID:	Q9QUN7
Synonyms:	Ly105



[View online »](#)

Summary:

Cooperates with LY96 to mediate the innate immune response to bacterial lipoproteins and other microbial cell wall components. Cooperates with TLR1 or TLR6 to mediate the innate immune response to bacterial lipoproteins or lipopeptides. Acts via MYD88 and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response (By similarity) (PubMed:15690042). May also promote apoptosis in response to lipoproteins (By similarity). Forms activation clusters composed of several receptors depending on the ligand, these clusters trigger signaling from the cell surface and subsequently are targeted to the Golgi in a lipid-raft dependent pathway. Forms the cluster TLR2:TLR6:CD14:CD36 in response to diacylated lipopeptides and TLR2:TLR1:CD14 in response to triacylated lipopeptides (By similarity). Recognizes M.tuberculosis major T-antigen EsxA (ESAT-6) which inhibits downstream MYD88-dependent signaling (PubMed:17486091). Acts as the major receptor for M.tuberculosis lipoproteins LprA, LprG, LpqH and PhoS1 (pstS1), in conjunction with TLR1 and for some but not all lipoproteins CD14 and/or CD36. The lipoproteins act as agonists to modulate antigen presenting cell functions in response to the pathogen (PubMed:19362712). Recombinant MPT83 from M.tuberculosis stimulates secretion of cytokines (TNF-alpha, IL-6 and IL-12p40) by mouse macrophage cell lines in a TLR2-dependent fashion, which leads to increased host innate immunity responses against the bacterium (PubMed:22174456). Lung macrophages which express low levels of TLR2 respond poorly to stimulation by M.tuberculosis LpqH (PubMed:19362712). Required for normal uptake of M.tuberculosis, a process that is inhibited by M.tuberculosis LppM (PubMed:27220037). Interacts with TICAM2 (By similarity).[UniProtKB/Swiss-Prot Function]

Product images: