

## Product datasheet for **KN318768BN**

### Usp13 Mouse Gene Knockout Kit (CRISPR)

#### Product data:

Product Type:	Knockout Kits (CRISPR)
Format:	2 gRNA vectors, 1 mBFP-Neo donor, 1 scramble control
Donor DNA:	mBFP-Neo
Symbol:	Usp13
Locus ID:	72607
Components:	<b>KN318768G1</b> , Usp13 gRNA vector 1 in pCas-Guide CRISPR vector (GE100002) <b>KN318768G2</b> , Usp13 gRNA vector 2 in pCas-Guide CRISPR vector (GE100002) <b>KN318768BND</b> , donor DNA containing left and right homologous arms and mBFP-Neo functional cassette. <b>GE100003</b> , scramble sequence in pCas-Guide vector
Disclaimer:	These products are manufactured and supplied by OriGene under license from ERS. The kit is designed based on the best knowledge of CRISPR technology. The system has been functionally validated for knocking-in the cassette downstream the native promoter. The efficiency of the knock-out varies due to the nature of the biology and the complexity of the experimental process.
RefSeq:	<a href="#">NM_001013024</a>
UniProt ID:	<a href="#">Q5BKP2</a>
Synonyms:	2700071E21Rik; AI848077; IsoT-3; ISOT3



[View online »](#)

**Summary:**

Deubiquitinase that mediates deubiquitination of target proteins such as BECN1, MITF, SKP2 and USP10 and is involved in various processes such as autophagy and endoplasmic reticulum-associated degradation (ERAD). Component of a regulatory loop that controls autophagy and p53/TP53 levels: mediates deubiquitination of BECN1, a key regulator of autophagy, leading to stabilize the PIK3C3/VPS34-containing complexes. Also deubiquitinates USP10, an essential regulator of p53/TP53 stability. In turn, PIK3C3/VPS34-containing complexes regulate USP13 stability, suggesting the existence of a regulatory system by which PIK3C3/VPS34-containing complexes regulate p53/TP53 protein levels via USP10 and USP13. Recruited by nuclear UFD1 and mediates deubiquitination of SKP2, thereby regulating endoplasmic reticulum-associated degradation (ERAD). Also regulates ERAD through the deubiquitination of UBL4A a component of the BAG6/BAT3 complex. Mediates stabilization of SIAH2 independently of deubiquitinase activity: binds ubiquitinated SIAH2 and acts by impairing SIAH2 autoubiquitination. Has a weak deubiquitinase activity in vitro and preferentially cleaves 'Lys-63'-linked polyubiquitin chains. In contrast to USP5, it is not able to mediate unanchored polyubiquitin disassembly. Able to cleave ISG15 in vitro; however, additional experiments are required to confirm such data.[UniProtKB/Swiss-Prot Function]

**Product images:**
