

## Product datasheet for **KN407772**

### FFAR3 Human Gene Knockout Kit (CRISPR)

#### Product data:

Product Type:	Knockout Kits (CRISPR)
Format:	2 gRNA vectors, 1 linear donor
Donor DNA:	EF1a-GFP-P2A-Puro
Symbol:	FFAR3
Locus ID:	2865



[View online »](#)

**Components:**
**KN407772G1**, FFAR3 gRNA vector 1 in pCas-Guide CRISPR vector (GE100002)

**KN407772G2**, FFAR3 gRNA vector 2 in pCas-Guide CRISPR vector (GE100002)

**KN407772D**, Linear donor DNA containing LoxP-EF1A-tGFP-P2A-Puro-LoxP:

The sequence below is cassette sequence only. The linear donor DNA also contains proprietary target sequence.

LoxP-EF1A-tGFP-P2A-Puro-LoxP (2739 bp)

```

ATAACTTCGT ATAATGTATG CTATACGAAG TTATCGTGAG GCTCCGGTGC CCGTCAGTGG GCAGAGCGCA
CATCGCCAC AGTCCCCGAG AAGTTGGGG GAGGGGTCCG CAATTGAACC GGTGCCTAGA GAAGGTGGCG
CGGGTAAC TGGGAAAGTG ATGTCGTGTA CTGGCTCCG CTTTTCCCG AGGGTGGGG AGAACCGTAT
ATAAGTCAG TAGTCGCCG GAACGTTCT TTTCCGAAC GGTTCGCCG CAGAACACAG GTAAGTGCCG
TGTGTGGTTC CCGCGGGCT GGCCTCTTA CGGGTTATG CCCTTGCGTG CCTTGAATTA CTTCCACCTG
GCTGCAGTAC GTGATTCTG ATCCCGAGCT TCGGGTTGGA AGTGGGTGG AGAGTTCGAG GCCTTGCGCT
TAAGGAGCCC CTTCGCCTG TGCTTGAGT GAGGCCTGCC CTGGGCGTG GGGCCCGCG GTGCGAATCT
GGTGGCACCT TCGCGCCTG CTCGCTGCT TCGATAAGT TCTAGCCATT TAAAATTTT GATGACCTGC
TGCAGCGCT TTTTCTGGC AAGATAGTCT TGTAATGCG GGCCAAGATC TGCACACTGG TATTTTCGTT
TTTGGGGCCG CGGGCGGCGA CGGGGCCCGT GCGTCCCAGC GCACATGTTC GGCAGGCGG GGCCTGCGAG
CGCGGCCACC GAGAATCGGA CGGGGTAGT CTCAAGCTGG CCGGCCTGCT CTGGTGCCTG GCCTCGCGCC
GCCGTGTATC GCCCGCCCT GGGCGGAAG GCTGGCCCG TCGGCACCAG TTGCGTGAGC GGAAAGATGG
CCGTTCCCG GCCCTGTGC AGGGAGCTCA AAATGGAGGA CGCGCGCTC GGGAGAGCGG GCGGGTGAAGT
CACCCACACA AAGGAAAAG GCCTTCCGT CCTCAGCCG CGCTTCATGT GACTCCACGG AGTACCGGGC
GCCGTCAGG CACCTCGAT AGTTCTCGAG CTTTGGAGT ACGTCGTCT TAGGTTGGGG GGAGGGGTTT
TATGCGATGG AGTTTCCCA CACTGAGTGG GTGAGACTG AAGTTAGGCC AGCTTGGCAG TTGATGTAAT
TCTCCTTGG AATTGCCCT TTTGAGTTG GATCTTGGT CATTCTCAAG CCTCAGACAG TGGTTCAAAG
TTTTTTCTT CCATTTCAAG TGTCGTGAAT GGAGAGCGAC GAGAGCGGCC TGCCCGCCAT GGAGATCGAG
TGCCGCATCA CCGGCACCCT GAACGGCGTG GAGTTCGAGC TGGTGGGCGG CGGAGAGGGC ACCCCCGAGC
AGGGCCGCAT GACCAACAAG ATGAAGAGCA CCAAAGGCGC CCTGACCTTC AGCCCTACC TGCTGAGCCA
CGTGATGGC TACGGCTTCT ACCACTTCG CACCTACCC AGCGGCTACG AGAACCCCTT CCTGCACGCC
ATCAACAACG GCGGCTACAC CAACACCCG ATCGAGAAGT ACGAGGACGG CGGCGTGCTG CACGTGAGCT
TCAGCTACCG CTACGAGGCC GGCCGCGTGA TCGGCGACTT CAAGGTGATG GGCACCGGCT TCCCGGAGGA
CAGCGTGATC TTCACCGACA AGATCATCCG CAGCAACGCC ACCGTGGAGC ACCTGCACCC CATGGCGGAT
AACGATCTGG ATGGCAGCTT CACCCGACC TTCAGCCTGC GCGACGGCGG CTACTIONAGC TCCGTGGTGG
ACAGCCACAT GCACTTCAAG AGCGCCATCC ACCCCAGCAT CCTGCAGAAC GGGGGCCCA TGTTCCCTT
CCGCCCGTG GAGGAGGATC ACAGCAACAC CGAGTGGGG ATCGTGGAGT ACCAGCACGC CTTCAAGACC
CCGGATGCAG ATGCCGGTGA AGAAAGAGGA AGCGGAGCTA CTAACCTCAG CCTGCTGAAG CAGGCTGGAG
ACGTGGAGGA GAACCTGGA CCTATGACCG AGTACAAGC CACGGTGC GCCTGCCACC GCGACGACGT
CCCCAGGGC GTACGCACC TCGCCGCCG GTTCGCCGAC TACCCGCCA CGGCCACAC CGTCGATCCG
GACCGCCACA TCGAGCGGT CACCGAGCTG CAAGAACTCT TCCTCACCG CGTCGGGCTC GACATCGGCA
AGGTGTGGT CGCGGACGAC GGCGCCGCG TGGCGTCTG GACCACGCC GAGAGCGTGC AAGCGGGGGC
GGTGTTCGCC GAGATCGGCC CGCGCATGGC CGAGTTGAGC GGTTCGGGC TGGCCCGCA GCAACAGATG
GAAGCCCTCC TGCGCCGCA CCGGCCAAG GAGCCCGCT GTTCTCTGG CACCCTCGG GTCTCGCCG
ACCACCAGG CAAGGTCTG GGCAGCGCG TCGTGTCCC CGGAGTGGAG GCGGCCGAGC GCGCCGGGT
GCCCGCTTC CTGGAGACT CCGCGCCCG CAACCTCCC TTCTACGAG GGCTCGGCT CACCGTCACC
GCCGACGTC AGGTGCCGA AGGACCGCG ACCTGGTGA TGACCCGCA GCCCGGTGCC TGAAACTTGT
TTATTGCAGC TTATAATGGT TACAAATAA GCAATAGCAT CACAAATTC ACAAATAAG CATTTTTTTC
ACTGCATTCT AGTTGTGGT TGTCCAACT CATCAATGA TCTTAATAAC TTCGTATAAT GTATGCTATA CGAAGTTAT
    
```



<b>Disclaimer:</b>	These products are manufactured and supplied by OriGene under license from ERS. The kit is designed based on the best knowledge of CRISPR technology. The system has been functionally validated for knocking-in the cassette downstream the native promoter. The efficiency of the knock-out varies due to the nature of the biology and the complexity of the experimental process.
<b>RefSeq:</b>	<a href="#">NM_005304</a>
<b>UniProt ID:</b>	<a href="#">O14843</a>
<b>Synonyms:</b>	FFA3R; GPR41
<b>Summary:</b>	<p>G protein-coupled receptor that is activated by a major product of dietary fiber digestion, the short chain fatty acids (SCFAs), and that plays a role in the regulation of whole-body energy homeostasis and in intestinal immunity. In omnivorous mammals, the short chain fatty acids acetate, propionate and butyrate are produced primarily by the gut microbiome that metabolizes dietary fibers. SCFAs serve as a source of energy but also act as signaling molecules. That G protein-coupled receptor is probably coupled to the pertussis toxin-sensitive, G(i/o)-alpha family of G proteins. Its activation results in the formation of inositol 1,4,5-trisphosphate, the mobilization of intracellular calcium, the phosphorylation of the MAPK3/ERK1 and MAPK1/ERK2 kinases and the inhibition of intracellular cAMP accumulation (PubMed:12711604). Activated by SCFAs and by beta-hydroxybutyrate, a ketone body produced by the liver upon starvation, it inhibits N-type calcium channels and modulates the activity of sympathetic neurons through a signaling cascade involving the beta and gamma subunits of its coupled G protein, phospholipase C and MAP kinases. Thereby, it may regulate energy expenditure through the control of the sympathetic nervous system that controls for instance heart rate. Upon activation by SCFAs accumulating in the intestine, it may also signal to the brain via neural circuits which in turn would regulate intestinal gluconeogenesis. May also control the production of hormones involved in whole-body energy homeostasis. May for instance, regulate blood pressure through renin secretion. May also regulate secretion of the PYY peptide by enteroendocrine cells and control gut motility, intestinal transit rate, and the harvesting of energy from SCFAs produced by gut microbiota. May also indirectly regulate the production of LEP/Leptin, a hormone acting on the CNS to inhibit food intake, in response to the presence of short-chain fatty acids in the intestine. Finally, may also play a role in glucose homeostasis. Besides its role in energy homeostasis, may play a role in intestinal immunity. May mediate the activation of the inflammatory and immune response by SCFAs in the gut, regulating the rapid production of chemokines and cytokines by intestinal epithelial cells. Among SCFAs, the fatty acids containing less than 6 carbons, the most potent activators are probably propionate, butyrate and pentanoate while acetate is a poor activator (PubMed:12496283, PubMed:12711604).[UniProtKB/Swiss-Prot Function]</p>

**Product images:**
