

Product datasheet for **KN413356**

TXK Human Gene Knockout Kit (CRISPR)

Product data:

Product Type:	Knockout Kits (CRISPR)
Format:	2 gRNA vectors, 1 linear donor
Donor DNA:	EF1a-GFP-P2A-Puro
Symbol:	TXK
Locus ID:	7294



[View online »](#)

Components:
KN413356G1, TXK gRNA vector 1 in pCas-Guide CRISPR vector (GE100002)

KN413356G2, TXK gRNA vector 2 in pCas-Guide CRISPR vector (GE100002)

KN413356D, Linear donor DNA containing LoxP-EF1a-tGFP-P2A-Puro-LoxP:

The sequence below is cassette sequence only. The linear donor DNA also contains proprietary target sequence.

LoxP-EF1a-tGFP-P2A-Puro-LoxP (2739 bp)

```

ATAACTTCGT ATAATGTATG CTATACGAAG TTATCGTGAG GCTCCGGTGC CCGTCAGTGG GCAGAGCGCA
CATCGCCAC AGTCCCGAG AAGTTGGGG GAGGGTTCGG CAATTGAACC GGTGCCTAGA GAAGGTGGCG
CGGGTAAC TGGAAAGTG ATGTCGTGTA CTGGCTCCG CTTTTCCCG AGGGTGGGG AGAACCGTAT
ATAAGTCAG TAGTCGCCG GAACGTTCT TTTCCGAAC GGTTCGCCG CAGAACACAG GTAAGTGCCG
TGTGTGGTTC CCGCGGGCT GGCCTCTTA CGGGTTATG CCCTTGCGTG CCTTGAATTA CTTCCACCTG
GCTGCAGTAC GTGATTCTG ATCCCGAGCT TCGGGTTGA AGTGGGTGG AGAGTTCGAG GCCTTGCGCT
TAAGGAGCCC CTTCGCCTG TGCTTGAGT GAGGCCTGC CTGGGCGTG GGGCCCGCG GTGCGAATCT
GGTGGCACCT TCGCGCCTG CTCGCTGCT TCGATAAGT TCTAGCCATT TAAAATTTT GATGACCTGC
TGCAGCGCT TTTTCTGGC AAGATAGTCT TGTAATGCG GGCCAAGATC TGCACACTGG TATTTTCGTT
TTTGGGGCCG CGGGCGGCA CGGGGCCCGT GCGTCCCAGC GCACATGTTC GGCAGGCGG GGCCTGCGAG
CGCGGCCACC GAGAATCGGA CGGGGTAGT CCAAGCTGG CCGGCCTGCT CTGGTGCCTG GCCTCGCGCC
GCCGTGTATC GCCCGCCCT GGGCGCAAG GCTGGCCCG TCGGCACCAG TTGCGTGAGC GGAAAGATGG
CCGTTCCCG GCCCTGTGC AGGGAGCTCA AAATGGAGGA CGCGCGCTC GGGAGAGCG GCGGGTGAAGT
CACCCACACA AAGGAAAAG GCCTTCCGT CCTCAGCCG CGCTTCATGT GACTCCACGG AGTACCGGGC
GCCGTCAGG CACCTCGAT AGTTCTCGAG CTTTGGAGT ACGTCGTCT TAGTTGGGG GGAGGGTTT
TATGCGATGG AGTTCCCCA CACTGAGTG GTGAGACTG AAGTTAGGCC AGCTTGGCAG TTGATGTAAT
TCTCCTTGA ATTTCCCTT TTTGAGTTG GATCTTGGT CATTCTCAAG CCTCAGACAG TGGTTCAAAG
TTTTTTCTT CCATTTCAAG TGTCGTGAAT GGAGAGCGAC GAGAGCGGCC TGCCCGCCAT GGAGATCGAG
TGCCGCATCA CCGGCACCCT GAACGGCGTG GAGTTCGAGC TGGTGGGCGG CGGAGAGGGC ACCCCGAGC
AGGGCCGCAT GACCAACAAG ATGAAGAGCA CCAAAGGCGC CCTGACCTTC AGCCCTACC TGCTGAGCCA
CGTGATGGC TACGGCTTCT ACCACTTCG CACCTACCC AGCGGCTACG AGAACCCCTT CCTGCACGCC
ATCAACAACG GCGGCTACAC CAACACCCG ATCGAGAAGT ACGAGGACGG CGGCGTGCTG CACGTGAGCT
TCAGTACCG CTACGAGGC GCGCGCTGA TCGGCGACTT CAAGGTGATG GGCACCGCT TCCCGGAGGA
CAGCGTGATC TTCACCGACA AGATCATCCG CAGCAACGCC ACCGTGGAGC ACCTGCACCC CATGGCGAT
AACGATCTGG ATGGCAGCTT CACCCGACC TTCAGCCTGC GCGACGGCGG CTACTACAGC TCCGTGGTGG
ACAGCCACAT GCACTTCAAG AGCGCCATCC ACCCCAGCAT CCTGCAGAAC GGGGGCCCA TGTTCCCTT
CCGCCCGTG GAGGAGGATC ACAGCAACAC CGAGTGGGC ATCGTGGAGT ACCAGCACGC CTTCAAGACC
CCGATGACG ATGCCGGTGA AGAAAGAGGA AGCGGAGCTA CTAACCTCAG CCTGCTGAAG CAGGCTGGAG
ACGTGGAGGA GAACCTGGA CCTATGACCG AGTACAAGC CACGGTGC GCCTGCCACC GCGACGACGT
CCCCAGGGC GTACGCACC TCGCCGCCG GTTCGCCGAC TACCCGCCA CGGCCACAC CGTCGATCCG
GACCGCCACA TCGAGCGGT CACCGAGCTG CAAGAATCT TCCTCACGG CGTCGGGTC GACATCGCA
AGGTGTGGT CGCGGACGAC GCGCCCGCG TGGCGTCTG GACCACGGC GAGAGCGTCG AAGCGGGGGC
GGTGTTCGC GAGATCGGC CGCGCATGG CGAGTTGAGC GGTTCGGC TGGCCCGCA GCAACAGATG
GAAGCCCTC TGCGCCGCA CCGGCCAAG GAGCCCGCT GTTCTCTGG CACCCTCGG GTCTCGCCG
ACCACCAGG CAAGGTCTG GGCAGCGCG TCGTGTCCC CGGAGTGGAG GCGGCCGAGC GCGCCGGGT
GCCCGCTT CTGGAGACT CCGCGCCCG CAACCTCCC TTCTACGAG GGCTCGGCT CACCGTACC
GCCGACGTC AGGTGCCGA AGGACCGCG ACCTGGTGA TGACCCGCA GCCCGGTGCC TGAAACTTGT
TTATTGCAG TTATAATGG TACAAATAA GCAATAGCAT CACAAATTC ACAAATAAG CATTTTTTT
ACTGCATTCT AGTTGTGGT TGTCCAACT CATCAATGA TCTTAATAA TTCGTATAAT GTATGTATA CGAAGTTAT
    
```



Disclaimer: These products are manufactured and supplied by OriGene under license from ERS. The kit is designed based on the best knowledge of CRISPR technology. The system has been functionally validated for knocking-in the cassette downstream the native promoter. The efficiency of the knock-out varies due to the nature of the biology and the complexity of the experimental process.

RefSeq: [NM_003328](#)

UniProt ID: [P42681](#)

Synonyms: BTKL; PSCTK5; PTK4; RLK; TKL

Summary: Non-receptor tyrosine kinase that plays a redundant role with ITK in regulation of the adaptive immune response. Regulates the development, function and differentiation of conventional T-cells and nonconventional NKT-cells. When antigen presenting cells (APC) activate T-cell receptor (TCR), a series of phosphorylation lead to the recruitment of TXK to the cell membrane, where it is phosphorylated at Tyr-420. Phosphorylation leads to TXK full activation. Contributes also to signaling from many receptors and participates in multiple downstream pathways, including regulation of the actin cytoskeleton. Like ITK, can phosphorylate PLCG1, leading to its localization in lipid rafts and activation, followed by subsequent cleavage of its substrates. In turn, the endoplasmic reticulum releases calcium in the cytoplasm and the nuclear activator of activated T-cells (NFAT) translocates into the nucleus to perform its transcriptional duty. With PARP1 and EEF1A1, TXK forms a complex that acts as a T-helper 1 (Th1) cell-specific transcription factor and binds the promoter of IFNG to directly regulate its transcription, and is thus involved importantly in Th1 cytokine production. Phosphorylates both PARP1 and EEF1A1. Phosphorylates also key sites in LCP2 leading to the up-regulation of Th1 preferred cytokine IL-2. Phosphorylates 'Tyr-201' of CTLA4 which leads to the association of PI-3 kinase with the CTLA4 receptor.[UniProtKB/Swiss-Prot Function]

Product images:

