

Product datasheet for **KN417658**

SYT10 Human Gene Knockout Kit (CRISPR)

Product data:

Product Type:	Knockout Kits (CRISPR)
Format:	2 gRNA vectors, 1 linear donor
Donor DNA:	EF1a-GFP-P2A-Puro
Symbol:	SYT10
Locus ID:	341359



[View online »](#)

Components:
KN417658G1, SYT10 gRNA vector 1 in pCas-Guide CRISPR vector (GE100002)

KN417658G2, SYT10 gRNA vector 2 in pCas-Guide CRISPR vector (GE100002)

KN417658D, Linear donor DNA containing LoxP-EF1A-tGFP-P2A-Puro-LoxP:

The sequence below is cassette sequence only. The linear donor DNA also contains proprietary target sequence.

LoxP-EF1A-tGFP-P2A-Puro-LoxP (2739 bp)

```

ATAACTTCGT ATAATGTATG CTATACGAAG TTATCGTGAG GCTCCGGTGC CCGTCAGTGG GCAGAGCGCA
CATCGCCAC AGTCCCGAG AAGTTGGGG GAGGGGTGCG CAATTGAACC GGTGCCTAGA GAAGGTGGCG
CGGGGTAAAC TGGGAAAGTG ATGTCGTGTA CTGGCTCCGC CTTTTCCCG AGGGTGGGG AGAACCGTAT
ATAAGTCAG TAGTCGCCGT GAACGTTCTT TTTGCAACG GGTTCGCCG CAGAACACAG GTAAGTGCCG
TGTGTGGTTC CCGCGGGCCT GGCCTCTTTA CGGGTTATGG CCCTTGCGTG CCTTGAATTA CTTCCACCTG
GCTGCAGTAC GTGATTCTTG ATCCCGAGCT TCGGGTTGGA AGTGGGTGGG AGAGTTCGAG GCCTTGCGCT
TAAGGAGCCC CTTCGCCTCG TGCTTGAGTT GAGGCCTGGC CTGGGCGCTG GGGCCGCCG GTGCGAATCT
GGTGGCACCT TCGCGCCTGT CTCGCTGCTT TCGATAAGTC TCTAGCCATT TAAAAATTTT GATGACCTGC
TGCAGCGCTT TTTTCTGGC AAGATAGTCT TGTAAATGCG GGCCAAGATC TGCACACTGG TATTTTCGTT
TTTGGGGCCG CGGGCGGCGA CGGGGCCCGT GCGTCCCAGC GCACATGTTC GGCAGGCGG GGCCTGCGAG
CGCGGCCACC GAGAATCGGA CGGGGGTAGT CTCAAGCTGG CCGGCCTGCT CTGGTGCCTG GCCTCGCGCC
GCCGTGTATC GCCCGGCCCT GGGCGGCAAG GCTGGCCCGG TCGGCACCAAG TTGCGTGAGC GGAAAGATGG
CCGCTTCCCG GCCCTGCTGC AGGGAGCTCA AAATGGAGGA CGCGGCGCTC GGGAGAGCGG GCGGGTGAGT
CACCCACACA AAGGAAAAGG GCCTTCCCGT CCTCAGCCGT CGCTTCATGT GACTCCACGG AGTACCGGGC
GCCGTCCAGG CACCTCGATT AGTTCTCGAG CTTTGGAGT ACCTCGTCTT TAGGTTGGGG GGAGGGGTTT
TATGCGATGG AGTTTCCCA CACTGAGTGG GTGGAGACTG AAGTTAGGCC AGCTTGGCAG TTGATGTAAT
TCTCCTTGGG ATTTGCCCTT TTTGAGTTG GATCTTGGTT CATTCTCAAG CCTCAGACAG TGGTTCAAAG
TTTTTTCTT CCATTTCAAG TGTCGTGAAT GGAGAGCGAC GAGAGCGGCC TGCCCGCCAT GGAGATCGAG
TGCCGCATCA CCGGCACCTT GAACGGCGTG GAGTTCGAGC TGGTGGGCGG CGGAGAGGGC ACCCCGAGC
AGGGCCGCAT GACCAACAAG ATGAAGAGCA CCAAAGGCGC CCTGACCTTC AGCCCTACC TGCTGAGCCA
CGTGATGGG TACGGTTCTT ACCACTTCGG CACCTACCCC AGCGGCTACG AGAACCCCTT CCTGCACGCC
ATCAACAACG GCGGCTACAC CAACACCCG ATCGAGAAGT ACGAGGACGG CGGCGTGCTG CACGTGAGCT
TCAGCTACCG CTACGAGGCC GGCCGCGTGA TCGGCGACTT CAAGGTGATG GGCACCGGCT TCCCCGAGGA
CAGCGTGATC TTCACCGACA AGATCATCCG CAGCAACGCC ACCGTGGAGC ACCTGCACCC CATGGCGGAT
AACGATCTGG ATGGCAGCTT CACCCGACCC TTCAGCCTGC GCGACGGCGG CTAATACAGC TCCGTGGTGG
ACAGCCACAT GCACTTCAAG AGCGCCATCC ACCCCAGCAT CCTGCAGAAC GGGGGCCCCA TGTTTCGCTT
CCGCCCGGTG GAGGAGGATC ACAGCAACAC CGAGCTGGGC ATCGTGAGT ACCAGCACGC CTTCAAGACC
CCGGATGCAG ATGCCGGTGA AGAAAGAGGA AGCGGAGCTA CTAAGTTTCA CCGTCTGAAG CAGGCTGGAG
ACGTGGAGGA GAACCTTGA CCTATGACCG AGTACAAGCC CACGGTGCGC CTCGCCACCC GCGACGACGT
CCCCAGGGCC GTACGCACCC TCGCCGCCGC GTTCGCCGAC TACCCGCCA CGGCCACAC CGTCGATCCG
GACCGCCACA TCGAGCGGGT CACCGAGCTG CAAGAACTCT TCCTCACGCG CGTCGGGCTC GACATCGGCA
AGGTGTGGGT CGCGGACGAC GCGCGCCGCG TGGCGGTCTG GACCACGCCG GAGAGCGTGC AAGCGGGGGC
GGTGTTCGCC GAGATCGGCC CGCGCATGGC CGAGTTGAGC GGTTCCCGG TGCCCGCGCA GCAACAGATG
GAAGGCCTCC TGGCGCCGCA CCGGCCCAAG GAGCCCGCGT GGTTCTTGGC CACCGTCGGC GTCTCGCCCG
ACCACAGGG CAAGGTCTG GGCAGCGCCG TCGTGCTCCC CGGAGTGGAG GCGGCCGAGC GCGCCGGGGT
GCCCGCCTTC CTGGAGACCT CCGCGCCCCG CAACCTCCCC TTCTACGAGC GGCTCGGCTT CACCGTCACC
GCCGACGTCG AGGTGCCGA AGGACCGCGC ACCTGGTGCA TGACCCGCAA GCCCGGTGCC TGAACTTGT
TTATTGCAGC TTATAATGGT TACAAATAAA GCAATAGCAT CACAAATTTT ACAAATAAAG CATTTTTTTC
ACTGCATTCT AGTTGTGGTT TGTCCAACT CATCAATGTA TCTTAATAAC TTCGTATAAT GTATGCTATA CGAAGTTAT
    
```



RefSeq:	<u>NM_198992</u>
UniProt ID:	<u>Q6XYQ8</u>
Synonyms:	MGC119436; MGC119437
Summary:	Ca(2+) sensor specifically required for the Ca(2+)-dependent exocytosis of secretory vesicles containing IGF1 in neurons of the olfactory bulb. Exocytosis of IGF1 is required for sensory perception of smell. Not involved in Ca(2+)-dependent synaptic vesicle exocytosis (By similarity). Acts through Ca(2+) and phospholipid binding to the C2 domain: Ca(2+) induces binding of the C2-domains to phospholipid membranes and to assembled SNARE-complexes; both actions contribute to triggering exocytosis (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:

