

Product datasheet for **KN418240**

PCDHB2 Human Gene Knockout Kit (CRISPR)

Product data:

Product Type:	Knockout Kits (CRISPR)
Format:	2 gRNA vectors, 1 linear donor
Donor DNA:	EF1a-GFP-P2A-Puro
Symbol:	PCDHB2
Locus ID:	56133



[View online »](#)

Components:
KN418240G1, PCDHB2 gRNA vector 1 in pCas-Guide CRISPR vector (GE100002)

KN418240G2, PCDHB2 gRNA vector 2 in pCas-Guide CRISPR vector (GE100002)

KN418240D, Linear donor DNA containing LoxP-EF1A-tGFP-P2A-Puro-LoxP:

The sequence below is cassette sequence only. The linear donor DNA also contains proprietary target sequence.

LoxP-EF1A-tGFP-P2A-Puro-LoxP (2739 bp)

ATAACTTCGT ATAATGTATG CTATACGAAG TTATCGTGAG GCTCCGGTGC CCGTCAGTGG GCAGAGCGCA
 CATCGCCAC AGTCCCGAG AAGTTGGGG GAGGGGTCGG CAATTGAACC GGTGCCTAGA GAAGGTGGCG
 CGGGGTAAAC TGGGAAAGTG ATGTCGTGTA CTGGCTCCGC CTTTTCCCG AGGGTGGGG AGAACCGTAT
 ATAAGTCAG TAGTCGCCGT GAACGTTCTT TTTGCAACG GGTTCGCCG CAGAACACAG GTAAGTGCCG
 TGTGTGGTTC CCGCGGGCCT GGCCTCTTTA CGGGTTATGG CCCTTGCGTG CCTTGAATTA CTTCCACCTG
 GCTGCAGTAC GTGATTCTTG ATCCGAGCT TCGGGTTGGA AGTGGGTGGG AGAGTTCGAG GCCTTGCGCT
 TAAGGAGCCC CTTCGCCTCG TGCTTGAGTT GAGGCCTGGC CTGGGCGCTG GGGCCGCCG GTGCGAATCT
 GGTGGCACCT TCGCGCCTGT CTCGCTGCTT TCGATAAGTC TCTAGCCATT TAAAATTTT GATGACCTGC
 TCGCAGCCTT TTTTCTGGC AAGATAGTCT TGTAAATGCG GGCCAAGATC TGCACACTGG TATTTTCGTT
 TTTGGGGCCG CGGGCGGCGA CGGGGCCCGT GCGTCCCAGC GCACATGTTC GGCAGGCGG GGCCTGCGAG
 CGCGGCCACC GAGAATCGGA CGGGGGTAGT CTAAGCTGG CCGGCCTGCT CTGGTGCCTG GCCTCGCGCC
 GCCGTGTATC GCCCGGCCCT GGGCGGCAAG GCTGGCCCG TCGGCACCAAG TTGCGTGAGC GGAAAGATGG
 CCGCTTCCCG GCCCTGCTGC AGGGAGCTCA AAATGGAGGA CGCGGCGCTC GGGAGAGCGG GCGGGTGAGT
 CACCCACACA AAGGAAAAGG GCCTTTCCGT CCTCAGCCGT CGCTTCATGT GACTCCACGG AGTACCGGGC
 GCCGTCCAGG CACCTCGATT AGTTCTCGAG CTTTGGAGT ACGTCGCTT TAGGTTGGGG GGAGGGGTTT
 TATGCGATGG AGTTTCCCCA CACTGAGTGG GTGGAGACTG AAGTTAGGCC AGCTTGGCAC TTGATGTAAT
 TCTCCTTGGG ATTTGCCCTT TTTGAGTTG GATCTTGGTT CATTCTCAAG CCTCAGACAG TGGTTCAAAG
 TTTTTTCTT CCATTTCAAG TGTCGTGAAT GGAGAGCGAC GAGAGCGGCC TGCCCGCCAT GGAGATCGAG
 TGCCGCATCA CCGGCACCTT GAACGGCGTG GAGTTCGAGC TGGTGGGCGG CGGAGAGGGC ACCCCGAGC
 AGGGCCGCAT GACCAACAAG ATGAAGAGCA CCAAAGGCGC CCTGACCTTC AGCCCTACC TGCTGAGCCA
 CGTGATGGG TACGGTTCTT ACCACTTCGG CACCTACCCC AGCGGCTACG AGAACCCCTT CCTGCACGCC
 ATCAACAACG GCGGCTACAC CAACACCCG ATCGAGAAGT ACGAGGACGG CGGCGTGCTG CACGTGAGCT
 TCAGCTACCG CTACGAGGCC GGCCGCGTGA TCGGCGACTT CAAGGTGATG GGCACCGGCT TCCCCGAGGA
 CAGCGTGATC TTCACCGACA AGATCATCCG CAGCAACGCC ACCGTGGAGC ACCTGCACCC CATGGCGGAT
 AACGATCTGG ATGGCAGCTT CACCCGACC TTCAGCCTGC GCGACGGCGG CTAATACAGC TCCGTGGTGG
 ACAGCCACAT GCACTTCAAG AGCGCCATCC ACCCCAGCAT CCTGCAGAAC GGGGGCCCCA TGTTCCGCTT
 CCGCCGCGTG GAGGAGGATC ACAGCAACAC CGAGCTGGGC ATCGTGAGT ACCAGCACGC CTTCAAGACC
 CCGGATGCAG ATGCCGGTGA AGAAAGAGGA AGCGGAGCTA CTAACCTCAG CCTGCTGAAG CAGGCTGGAG
 ACGTGGAGGA GAACCTTGA CCTATGACCG AGTACAAGCC CACGGTGCGC CTCGCCACCC GCGACGACGT
 CCCAGGGCC GTACGCACCC TCGCCGCCG GTTCGCCGAC TACCCGCA CGGCCACAC CGTCGATCCG
 GACCGCCACA TCGAGCGGT CACCGAGCTG CAAGAACTCT TCCTCACGCG CGTCGGGCTC GACATCGGCA
 AGGTGTGGT GCGGACGAC GCGCCCGCG TGGCGGTCTG GACCACGCC GAGAGCGTCG AAGCGGGGGC
 GGTGTTCCGC GAGATCGGCC CGCGCATGGC CGAGTTGAGC GGTTCGCGG TGCCCGCGCA GCAACAGATG
 GAAGGCCCTC TGCGCCGCA CCGGCCAAG GAGCCCGCT GGTTCCTGGC CACCGTCGGC GTCTCGCCG
 ACCACAGGG CAAGGTCTG GGCAGCGCG TCGTGCTCC CGAGTGAGG GCGGCCGAGC GCGCCGGGT
 GCCCGCTTC CTGGAGACCT CCGCGCCCC CAACCTCCC TTCTACGAG GGCTCGGCTT CACCGTCACC
 GCCGACGTC AGGTGCCGA AGGACCGCG ACCTGGTGCA TGACCCGCAA GCCCGGTGCC TGAACTTGT
 TTATTGCAGC TTATAATGGT TACAAATAA GCAATAGCAT CACAAATTTT ACAAATAAAG CATTTTTTTC
 ACTGCATTCT AGTTGTGGT TGTCCAACT CATCAATGTA TCTTAATAA TTCGTATAAT GTATGCTATA CGAAGTTAT



RefSeq: [NM_018936](#)

UniProt ID: [Q9Y5E7](#)

Synonyms: PCDH-BETA2

Summary: This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike the alpha and gamma clusters, the transcripts from these genes are made up of only one large exon, not sharing common 3' exons as expected. These neural cadherin-like cell adhesion proteins are integral plasma membrane proteins. Their specific functions are unknown but they most likely play a critical role in the establishment and function of specific cell-cell neural connections. [provided by RefSeq, Jul 2008]

Product images:

