

Product datasheet for **KN500317**

Cyren Mouse Gene Knockout Kit (CRISPR)

Product data:

Product Type:	Knockout Kits (CRISPR)
Format:	2 gRNA vectors, 1 linear donor
Donor DNA:	EF1a-GFP-P2A-Puro
Symbol:	Cyren
Locus ID:	78412



[View online »](#)

Components:
KN500317G1, Cyren gRNA vector 1 in pCas-Guide CRISPR vector (GE100002)

KN500317G2, Cyren gRNA vector 2 in pCas-Guide CRISPR vector (GE100002)

KN500317D, Linear donor DNA containing LoxP-EF1A-tGFP-P2A-Puro-LoxP:

The sequence below is cassette sequence only. The linear donor DNA also contains proprietary target sequence.

LoxP-EF1A-tGFP-P2A-Puro-LoxP (2739 bp)

```

ATAACTTCGT ATAATGTATG CTATACGAAG TTATCGTGAG GCTCCGGTGC CCGTCAGTGG GCAGAGCGCA
CATCGCCAC AGTCCCGAG AAGTTGGGG GAGGGGTGCG CAATTGAACC GGTGCCTAGA GAAGGTGGCG
CGGGGTAAAC TGGGAAAGTG ATGTCGTGTA CTGGCTCCGC CTTTTTCCCG AGGGTGGGG AGAACCGTAT
ATAAGTCAG TAGTCGCCGT GAACGTTCTT TTTGCAACG GGTTCGCCG CAGAACACAG GTAAGTGCCG
TGTGTGGTTC CCGCGGGCCT GGCCTCTTTA CGGGTTATGG CCCTTGCGTG CCTTGAATTA CTTCCACCTG
GCTGCAGTAC GTGATTCTTG ATCCCGAGCT TCGGGTTGGA AGTGGGTGGG AGAGTTCGAG GCCTTGCGCT
TAAGGAGCCC CTTCGCCTCG TGCTTGAGTT GAGGCCTGGC CTGGGCGCTG GGGCCGCCG GTGCGAATCT
GGTGGCACCT TCGCGCCTGT CTCGCTGCTT TCGATAAGTC TCTAGCCATT TAAAATTTT GATGACCTGC
TGCAGCGCTT TTTTCTGGC AAGATAGTCT TGTAAATGCG GGCCAAGATC TGCACACTGG TATTTTCGTT
TTTGGGGCCG CGGGCGGCGA CGGGGCCCGT GCGTCCCAGC GCACATGTTC GGCAGGCGG GGCCTGCGAG
CGCGGCCACC GAGAATCGGA CGGGGGTAGT CTCAAGCTGG CCGGCCTGCT CTGGTGCCTG GCCTCGCGCC
GCCGTGTATC GCCCGGCCCT GGGCGGCAAG GCTGGCCCGG TCGGCACCAAG TTGCGTGAGC GGAAAGATGG
CCGCTTCCCG GCCCTGCTGC AGGGAGCTCA AAATGGAGGA CGCGGCGCTC GGGAGAGCGG GCGGGTGAGT
CACCCACACA AAGGAAAAGG GCCTTTCCTG CCTCAGCCGT CGCTTCATGT GACTCCACGG AGTACCGGGC
GCCGTCCAGG CACCTCGATT AGTTCTCGAG CTTTGGAGT ACGTCGCTT TAGGTTGGGG GGAGGGGTTT
TATGCGATGG AGTTTCCCA CACTGAGTGG GTGGAGACTG AAGTTAGGCC AGCTTGGCAG TTGATGTAAT
TCTCCTTGGG ATTTGCCCTT TTTGAGTTTG GATCTTGGTT CATTCTCAAG CCTCAGACAG TGGTTCAAAG
TTTTTTCTT CCATTTCAAG TGTCGTGAAT GGAGAGCGAC GAGAGCGGCC TGCCCGCCAT GGAGATCGAG
TGCCGCATCA CCGGCACCTT GAACGGCGTG GAGTTCGAGC TGGTGGGCGG CGGAGAGGGC ACCCCGAGC
AGGGCCGCAT GACCAACAAG ATGAAGAGCA CCAAAGGCGC CCTGACCTTC AGCCCTACC TGCTGAGCCA
CGTGATGGG TACGGTTCTT ACCACTTCGG CACCTACCCC AGCGGCTACG AGAACCCCTT CCTGCACGCC
ATCAACAACG GCGGCTACAC CAACACCCG ATCGAGAAGT ACGAGGACGG CGGCGTGCTG CACGTGAGCT
TCAGCTACCG CTACGAGGCC GGCCGCGTGA TCGGCGACTT CAAGGTGATG GGCACCGGCT TCCCCGAGGA
CAGCGTGATC TTCACCGACA AGATCATCCG CAGCAACGCC ACCGTGGAGC ACCTGCACCC CATGGCGGAT
AACGATCTGG ATGGCAGCTT CACCCGACCC TTCAGCCTGC GCGACGGCGG CTAATACAGC TCCGTGGTGG
ACAGCCACAT GCACTTCAAG AGCGCCATCC ACCCCAGCAT CCTGCAGAAC GGGGGCCCCA TGTTTCGCTT
CCGCCCGGTG GAGGAGGATC ACAGCAACAC CGAGCTGGGC ATCGTGAGT ACCAGCACGC CTTCAAGACC
CCGGATGCAG ATGCCGGTGA AGAAAGAGGA AGCGGAGCTA CTAAGTTCAG CCTGCTGAAG CAGGCTGGAG
ACGTGGAGGA GAACCTTGA CCTATGACCG AGTACAAGCC CACGGTGCGC CTCGCCACCC GCGACGACGT
CCCCAGGGCC GTACGCACCC TCGCCGCCGC GTTCGCCGAC TACCCGCA CGGCCACAC CGTCGATCCG
GACCGCCACA TCGAGCGGGT CACCGAGCTG CAAGAACTCT TCCTCACGCG CGTCGGGCTC GACATCGGCA
AGGTGTGGT GCGGACGAC GCGCCCGCGG TGGCGGTCTG GACCACGCCG GAGAGCGTGC AAGCGGGGGC
GGTGTTCGCC GAGATCGGCC CGCGCATGGC CGAGTTGAGC GGTTCGCGG TGCCCGCGCA GCAACAGATG
GAAGGCCTCC TGGCGCCGCA CCGGCCAAG GAGCCCGCGT GGTTCCTGGC CACCGTCGGC GTCTCGCCCG
ACCACAGGG CAAGGTCTG GGCAGCGCCG TCGTGCTCC CGAGTGAGG GCGGCCGAGC GCGCCGGGT
GCCCGCTTC CTGGAGACCT CCGCGCCCCG CAACCTCCCC TTCTACGAG GGCTCGGCTT CACCGTCACC
GCCGACGTC AGGTGCCGA AGGACCGCGC ACCTGGTGCA TGACCCGCAA GCCCGGTGCC TGAACTTGT
TTATTGCAGC TTATAATGGT TACAAATAAA GCAATAGCAT CACAAATTTT ACAAATAAAG CATTTTTTTC
ACTGCATTCT AGTTGTGGT TGTCCAACT CATCAATGTA TCTTAATAA TTCGTATAAT GTATGCTATA CGAAGTTAT
    
```



RefSeq:	NM_001135611 , NM_001347101 , NM_199145
UniProt ID:	Q8BHZ5
Synonyms:	AL023049; Mri
Summary:	Cell-cycle-specific inhibitor of classical non-homologous end joining (NHEJ) of DNA double-strand break (DSB) repair during the S and G2 phases. Acts as a regulator of DNA repair pathway choice by specifically inhibiting classical NHEJ during the S and G2 phases, thereby promoting error-free repair by homologous recombination during cell cycle phases when sister chromatids are present. Preferentially protects single-stranded overhangs at break sites by inhibiting classical NHEJ, thereby creating a local environment that favors homologous recombination. Acts via interaction with XRCC5/Ku80 and XRCC6/Ku70, interaction restricted during the S and G2 phases only. Molecular mechanisms governing classical NHEJ inhibition via interaction with XRCC5/Ku80 and XRCC6/Ku70 are unknown (By similarity). May act as a regulator of proteasome (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:

