

Product datasheet for **KN505111**

Eif3g Mouse Gene Knockout Kit (CRISPR)

Product data:

Product Type:	Knockout Kits (CRISPR)
Format:	2 gRNA vectors, 1 linear donor
Donor DNA:	EF1a-GFP-P2A-Puro
Symbol:	Eif3g
Locus ID:	53356



[View online »](#)

Components:

KN505111G1, Eif3g gRNA vector 1 in pCas-Guide CRISPR vector (GE100002)

KN505111G2, Eif3g gRNA vector 2 in pCas-Guide CRISPR vector (GE100002)

KN505111D, Linear donor DNA containing LoxP-EF1A-tGFP-P2A-Puro-LoxP:

The sequence below is cassette sequence only. The linear donor DNA also contains proprietary target sequence.

LoxP-EF1A-tGFP-P2A-Puro-LoxP (2739 bp)

```

ATAACTTCGT ATAATGTATG CTATACGAAG TTATCGTGAG GCTCCGGTGC CCGTCAGTGG GCAGAGCGCA
CATCGCCAC AGTCCCGAG AAGTTGGGG GAGGGGTGCG CAATTGAACC GGTGCCTAGA GAAGGTGGCG
CGGGGTAAAC TGGGAAAGTG ATGTCGTGTA CTGGCTCCGC CTTTTCCCG AGGGTGGGG AGAACCGTAT
ATAAGTCAG TAGTCGCCGT GAACGTTCTT TTTGCAACG GGTTCGCCG CAGAACACAG GTAAGTGCCG
TGTGTGGTTC CCGCGGGCCT GGCCTCTTTA CGGGTTATGG CCCTTGCGTG CCTTGAATTA CTTCCACCTG
GCTGCAGTAC GTGATTCTTG ATCCGAGCT TCGGGTTGGA AGTGGGTGGG AGAGTTCGAG GCCTTGCGCT
TAAGGAGCCC CTTCGCCTCG TGCTTGAGTT GAGGCCTGGC CTGGGCGCTG GGGCCGCCG GTGCGAATCT
GGTGGCACCT TCGCGCCTGT CTCGCTGCTT TCGATAAGTC TCTAGCCATT TAAAAATTTT GATGACCTGC
TGCAGCGCTT TTTTCTGGC AAGATAGTCT TGTAAATGCG GGCCAAGATC TGCACACTGG TATTTCCGTT
TTTGGGGCCG CGGGCGGCGA CGGGGCCCGT GCGTCCCAGC GCACATGTTC GGCAGGCGG GGCCTGCGAG
CGCGGCCACC GAGAATCGGA CGGGGGTAGT CTCAAGCTGG CCGGCCTGCT CTGGTGCCTG GCCTCGCGCC
GCCGTGTATC GCCCGGCCCT GGGCGGCAAG GCTGGCCCGG TCGGCACCAAG TTGCGTGAGC GGAAAGATGG
CCGCTTCCCG GCCCTGCTGC AGGGAGCTCA AAATGGAGGA CGCGGCGCTC GGGAGAGCGG GCGGGTGAGT
CACCCACACA AAGGAAAAGG GCCTTCCCGT CCTCAGCCGT CGCTTCATGT GACTCCACGG AGTACCGGGC
GCCGTCCAGG CACCTCGATT AGTTCTCGAG CTTTGGAGT ACCTCGTCTT TAGGTTGGGG GGAGGGGTTT
TATGCGATGG AGTTTCCCA CACTGAGTGG GTGGAGACTG AAGTTAGGCC AGCTTGGCAG TTGATGTAAT
TCTCCTTGGG ATTTGCCCTT TTTGAGTTG GATCTTGGTT CATTCTCAAG CCTCAGACAG TGGTTCAAAG
TTTTTTCTT CCATTTCAAG TGTCGTGAAT GGAGAGCGAC GAGAGCGGCC TGCCCGCCAT GGAGATCGAG
TGCCGCATCA CCGGCACCTT GAACGGCGTG GAGTTCGAGC TGGTGGGCGG CGGAGAGGGC ACCCCGAGC
AGGGCCGCAT GACCAACAAG ATGAAGAGCA CCAAAGGCGC CCTGACCTTC AGCCCTACC TGCTGAGCCA
CGTGATGGG TACGGCTTCT ACCACTTCGG CACCTACCCC AGCGGCTACG AGAACCCCTT CCTGCACGCC
ATCAACAACG GCGGCTACAC CAACACCCG ATCGAGAAGT ACGAGGACGG CGGCGTGCTG CACGTGAGCT
TCAGCTACCG CTACGAGGCC GGCCGCGTGA TCGGCGACTT CAAGGTGATG GGCACCGGCT TCCCCGAGGA
CAGCGTGATC TTCACCGACA AGATCATCCG CAGCAACGCC ACCGTGGAGC ACCTGCACCC CATGGCGGAT
AACGATCTGG ATGGCAGCTT CACCCGACCC TTCAGCCTGC GCGACGGCGG CTAATACAGC TCCGTGGTGG
ACAGCCACAT GCACTTCAAG AGCGCCATCC ACCCCAGCAT CCTGCAGAAC GGGGGCCCCA TGTTCCGCTT
CCGCCCGGTG GAGGAGGATC ACAGCAACAC CGAGCTGGGC ATCGTGAGT ACCAGCACGC CTTCAAGACC
CCGGATGCAG ATGCCGGTGA AGAAAGAGGA AGCGGAGCTA CTAAGTTCAG CCTGCTGAAG CAGGCTGGAG
ACGTGGAGGA GAACCTTGA CCTATGACCG AGTACAAGCC CACGGTGCGC CTCGCCACCC GCGACGACGT
CCCCAGGGCC GTACGCACCC TCGCCGCCG GTTCGCCGAC TACCCGCCA CGGCCACAC CGTCGATCCG
GACCGCCACA TCGAGCGGGT CACCGAGCTG CAAGAACTCT TCCTCACGCG CGTCGGGCTC GACATCGGCA
AGGTGTGGT GCGGACGAC GCGCCCGCG TGGCGGTCTG GACCACGCC GAGAGCGTGC AAGCGGGGGC
GGTGTTCGCC GAGATCGGCC CGCGCATGGC CGAGTTGAGC GGTTCGCCG TGCCCGCGCA GCAACAGATG
GAAGGCCTCC TGGCGCCGCA CCGGCCAAG GAGCCCGCT GGTTCCTGGC CACCGTCGGC GTCTCGCCG
ACCACAGGG CAAGGTCTG GGCAGCGCC TCGTGCTCC CGAGTGAGG GCGGCCGAGC GCGCCGGGT
GCCCGCCTT CTGGAGACCT CCGCGCCCC CAACCTCCC TTCTACGAG GGCTCGGCTT CACCGTCACC
GCCGACGTC AGGTGCCGA AGGACCGCG ACCTGGTGCA TGACCCGCAA GCCCGGTGCC TGAACTTGT
TTATTGCAGC TTATAATGGT TACAAATAA GCAATAGCAT CACAAATTTT ACAAATAAAG CATTTTTTTC
ACTGCATTCT AGTTGTGGT TGTCCAACT CATCAATGTA TCTTAATAA TTCGTATAAT GTATGCTATA CGAAGTTAT
  
```



RefSeq:	NM_016876
UniProt ID:	Q9Z1D1
Synonyms:	44kDa; D0Jmb4; eIF3-p44; Eif3s4; p44; TU-189B2
Summary:	RNA-binding component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAⁱ and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression. This subunit can bind 18S rRNA. [UniProtKB/Swiss-Prot Function]

Product images:

