

Product datasheet for **KN508608**

Kcna2 Mouse Gene Knockout Kit (CRISPR)

Product data:

Product Type:	Knockout Kits (CRISPR)
Format:	2 gRNA vectors, 1 linear donor
Donor DNA:	EF1a-GFP-P2A-Puro
Symbol:	Kcna2
Locus ID:	16490



[View online »](#)

Components:
KN508608G1, Kcna2 gRNA vector 1 in pCas-Guide CRISPR vector (GE100002)

KN508608G2, Kcna2 gRNA vector 2 in pCas-Guide CRISPR vector (GE100002)

KN508608D, Linear donor DNA containing LoxP-EF1a-tGFP-P2A-Puro-LoxP:

The sequence below is cassette sequence only. The linear donor DNA also contains proprietary target sequence.

LoxP-EF1a-tGFP-P2A-Puro-LoxP (2739 bp)

```

ATAACTTCGT ATAATGTATG CTATACGAAG TTATCGTGAG GCTCCGGTGC CCGTCAGTGG GCAGAGCGCA
CATCGCCAC AGTCCCGAG AAGTTGGGG GAGGGTTCGG CAATTGAACC GGTGCCTAGA GAAGGTGGCG
CGGGTAAC TGGAAAGTG ATGTCGTGTA CTGGCTCCG CTTTTCCCG AGGGTGGGG AGAACCGTAT
ATAAGTCAG TAGTCGCCG GAACGTTCT TTTCCGAAC GGTTCGCCG CAGAACACAG GTAAGTGCCG
TGTGTGGTTC CCGCGGGCT GGCCTCTTA CGGGTTATG CCCTTGCGTG CCTTGAATTA CTTCCACCTG
GCTGCAGTAC GTGATTCTG ATCCCGAGCT TCGGGTTGA AGTGGGTGG AGAGTTCGAG GCCTTGCGCT
TAAGGAGCCC CTTCGCCTG TGCTTGAGT GAGGCCTGG CTGGGCGTG GGGCCCGCG GTGCGAATCT
GGTGGCACCT TCGCGCCTG CTCGCTGCT TCGATAAGT TCTAGCCATT TAAAATTTT GATGACCTGC
TGCAGCGCT TTTTCTGGC AAGATAGTCT TGTAATGCG GGCCAAGATC TGCACACTGG TATTTTCGTT
TTTGGGGCCG CGGGCGGCA CGGGGCCCGT GCGTCCCAGC GCACATGTTC GGCAGGCGG GGCCTGCGAG
CGCGGCCACC GAGAATCGGA CGGGGTAGT CTAAGCTGG CCGGCCTGCT CTGGTGCCTG GCCTCGCGCC
GCCGTGTATC GCCCGCCCT GGGCGGAAG GCTGGCCCG TCGGCACCAG TTGCGTGAGC GGAAAGATGG
CCGTTCCCG GCCCTGTGC AGGGAGCTCA AAATGGAGGA CGCGGCGCTC GGGAGAGCGG GCGGGTGAAGT
CACCCACACA AAGGAAAAGG GCCTTCCGT CCTCAGCCG CGCTTCATGT GACTCCACGG AGTACCGGGC
GCCGTCAGG CACCTCGAT AGTTCTCGAG CTTTTGAGT ACGTGCTCT TAGTTGGGG GGAGGGGTTT
TATGCGATGG AGTTTCCCA CACTGAGTGG GTGAGACTG AAGTTAGGCC AGCTTGGCAG TTGATGTAAT
TCTCCTTGG AATTGCCCT TTTGAGTTG GATCTTGGT CATTCTCAAG CCTCAGACAG TGGTTCAAAG
TTTTTTCTT CCATTTCAAG TGTCGTGAAT GGAGAGCGAC GAGAGCGGCC TGCCCGCCAT GGAGATCGAG
TGCCGCATCA CCGGCACCCT GAACGGCGTG GAGTTCGAGC TGGTGGGCGG CGGAGAGGGC ACCCCGAGC
AGGGCCGCAT GACCAACAAG ATGAAGAGCA CCAAAGGCGC CCTGACCTTC AGCCCTACC TGCTGAGCCA
CGTGATGGC TACGGCTTCT ACCACTTCG CACCTACCC AGCGGCTACG AGAACCCCTT CCTGCACGCC
ATCAACAACG GCGGCTACAC CAACACCCG ATCGAGAAGT ACGAGGACGG CGGCGTGCTG CACGTGAGCT
TCAGCTACCG CTACGAGGC GCGCGCTGA TCGGCGACTT CAAGGTGATG GGCACCGGCT TCCCGGAGGA
CAGCGTGATC TTCACCGACA AGATCATCCG CAGCAACGCC ACCGTGGAGC ACCTGCACCC CATGGCGGAT
AACGATCTGG ATGGCAGCTT CACCCGACC TTCAGCCTGC GCGACGGCGG CTACTACAGC TCCGTGGTGG
ACAGCCACAT GCACTTCAAG AGCGCCATCC ACCCCAGCAT CCTGCAGAAC GGGGGCCCA TGTTCCCTT
CCGCCCGTG GAGGAGGATC ACAGCAACAC CGAGTGGGC ATCGTGGAGT ACCAGCACGC CTTCAAGACC
CCGGATGCAG ATGCCGGTGA AGAAAGAGGA AGCGGAGCTA CTAACCTCAG CCTGCTGAAG CAGGCTGGAG
ACGTGGAGGA GAACCTGGA CCTATGACCG AGTACAAGC CACGGTGC GCCTGCCACC GCGACGACGT
CCCCAGGGC GTACGCACC TCGCCGCCG GTTCGCCGAC TACCCGCCA CGGCCACAC CGTCGATCCG
GACCGCCACA TCGAGCGGT CACCGAGCTG CAAGAAGCTT TCCTCACGG CGTCGGGCTC GACATCGGCA
AGGTGTGGT CCGGACGAC GCGCCCGCG TGGCGTCTG GACCACGCG GAGAGCGTCG AAGCGGGGGC
GGTGTTCGC GAGATCGGC CGCGCATGG CGAGTTGAGC GGTTCGGC TGGCCCGCA GCAACAGATG
GAAGCCCTC TGCGCCGCA CCGGCCAAG GAGCCCGCT GTTCTCTGG CACCCTCGG GTCTCGCCG
ACCACCAGG CAAGGTCTG GGCAGCGCG TCGTGTCCC CGGAGTGGAG GCGGCCGAGC GCGCCGGGT
GCCCGCTT CTGGAGACT CCGCGCCCG CAACCTCCC TTCTACGAG GGCTCGGCT CACCGTCACC
GCCGACGTC AGGTGCCGA AGGACCGCG ACCTGGTGA TGACCCGCA GCCCGGTGCC TGAACTTGT
TTATTGCAG TTATAATGG TACAAATAA GCAATAGCAT CACAAATTC ACAAATAAG CATTTTTTT
ACTGCATTCT AGTTGTGGT TGTCAAAC CATCAATGA TCTTAATAA TTCGTATAAT GTATGCTATA CGAAGTTAT
    
```



Disclaimer: These products are manufactured and supplied by OriGene under license from ERS. The kit is designed based on the best knowledge of CRISPR technology. The system has been functionally validated for knocking-in the cassette downstream the native promoter. The efficiency of the knock-out varies due to the nature of the biology and the complexity of the experimental process.

RefSeq: [NM_008417](#)

UniProt ID: [P63141](#)

Synonyms: Akr6a4; ENSMUSG00000074335; Gm10672; Kca1-2; Kv1.2; Mk-2

Summary:

Voltage-gated potassium channel that mediates transmembrane potassium transport in excitable membranes, primarily in the brain and the central nervous system, but also in the cardiovascular system. Prevents aberrant action potential firing and regulates neuronal output. Forms tetrameric potassium-selective channels through which potassium ions pass in accordance with their electrochemical gradient. The channel alternates between opened and closed conformations in response to the voltage difference across the membrane (PubMed:12527813, PubMed:21233214). Can form functional homotetrameric channels and heterotetrameric channels that contain variable proportions of KCNA1, KCNA2, KCNA4, KCNA5, KCNA6, KCNA7, and possibly other family members as well; channel properties depend on the type of alpha subunits that are part of the channel (PubMed:20696761). Channel properties are modulated by cytoplasmic beta subunits that regulate the subcellular location of the alpha subunits and promote rapid inactivation of delayed rectifier potassium channels (By similarity). In vivo, membranes probably contain a mixture of heteromeric potassium channel complexes, making it difficult to assign currents observed in intact tissues to any particular potassium channel family member. Homotetrameric KCNA2 forms a delayed-rectifier potassium channel that opens in response to membrane depolarization, followed by slow spontaneous channel closure (PubMed:23864368). In contrast, a heteromultimer formed by KCNA2 and KCNA4 shows rapid inactivation (PubMed:23864368). Contributes to the regulation of action potentials in neurons (PubMed:12527813, PubMed:17925011). KCNA2-containing channels play a presynaptic role and prevent hyperexcitability and aberrant action potential firing (PubMed:17634333, PubMed:17925011). Response to toxins that are selective for KCNA1, respectively for KCNA2, suggests that heteromeric potassium channels composed of both KCNA1 and KCNA2 play a role in pacemaking and regulate the output of deep cerebellar nuclear neurons (By similarity). Response to toxins that are selective for KCNA2-containing potassium channels suggests that in Purkinje cells, dendritic subthreshold KCNA2-containing potassium channels prevent random spontaneous calcium spikes, suppressing dendritic hyperexcitability without hindering the generation of somatic action potentials, and thereby play an important role in motor coordination (By similarity). KCNA2-containing channels play a role in GABAergic transmission from basket cells to Purkinje cells in the cerebellum, and thereby play an important role in motor coordination (PubMed:20696761). Plays a role in the induction of long-term potentiation of neuron excitability in the CA3 layer of the hippocampus (PubMed:23981714). May function as down-stream effector for G protein-coupled receptors and inhibit GABAergic inputs to basolateral amygdala neurons (By similarity). May contribute to the regulation of neurotransmitter release, such as gamma-aminobutyric acid (GABA) (By similarity). Contributes to the regulation of the axonal release of the neurotransmitter dopamine (PubMed:21233214). Reduced KCNA2 expression plays a role in the perception of neuropathic pain after peripheral nerve injury, but not acute pain (By similarity). Plays a role in the regulation of the time spent in non-rapid eye movement (NREM) sleep (PubMed:17925011).[UniProtKB/Swiss-Prot Function]

Product images:

