

Product datasheet for **KN510526**

Mug1 Mouse Gene Knockout Kit (CRISPR)

Product data:

Product Type:	Knockout Kits (CRISPR)
Format:	2 gRNA vectors, 1 linear donor
Donor DNA:	EF1a-GFP-P2A-Puro
Symbol:	Mug1
Locus ID:	17836



[View online »](#)

Components:
KN510526G1, Mug1 gRNA vector 1 in pCas-Guide CRISPR vector (GE100002)

KN510526G2, Mug1 gRNA vector 2 in pCas-Guide CRISPR vector (GE100002)

KN510526D, Linear donor DNA containing LoxP-EF1A-tGFP-P2A-Puro-LoxP:

The sequence below is cassette sequence only. The linear donor DNA also contains proprietary target sequence.

LoxP-EF1A-tGFP-P2A-Puro-LoxP (2739 bp)

```

ATAACTTCGT ATAATGTATG CTATACGAAG TTATCGTGAG GCTCCGGTGC CCGTCAGTGG GCAGAGCGCA
CATCGCCAC AGTCCCGAG AAGTTGGGG GAGGGGTGCG CAATTGAACC GGTGCCTAGA GAAGGTGGCG
CGGGGTAAAC TGGGAAAGTG ATGTCGTGTA CTGGCTCCGC CTTTTCCCG AGGGTGGGG AGAACCGTAT
ATAAGTCAG TAGTCGCCGT GAACGTTCTT TTTGCAACG GGTTCGCCG CAGAACACAG GTAAGTGCCG
TGTGTGGTTC CCGCGGGCCT GGCCTCTTTA CGGGTTATGG CCCTTGCGTG CCTTGAATTA CTTCCACCTG
GCTGCAGTAC GTGATTCTTG ATCCGAGCT TCGGGTTGGA AGTGGGTGGG AGAGTTCGAG GCCTTGCCTG
TAAGGAGCCC CTTCGCCTCG TGCTTGAGTT GAGGCCTGGC CTGGGCGCTG GGGCCGCCG GTGCGAATCT
GGTGGCACCT TCGCGCCTGT CTCGCTGCTT TCGATAAGTC TCTAGCCATT TAAAATTTT GATGACCTGC
TGCAGCGCTT TTTTCTGGC AAGATAGTCT TGTAAATGCG GGCCAAGATC TGCACACTGG TATTTCCGTT
TTTGGGGCCG CGGGCGGCGA CGGGGCCCGT GCGTCCCAGC GCACATGTTC GGCAGGCGG GGCCTGCGAG
CGCGGCCACC GAGAATCGGA CGGGGGTAGT CTCAAGCTGG CCGGCCTGCT CTGGTGCCTG GCCTCGCGCC
GCCGTGTATC GCCCGGCCCT GGGCGGCAAG GCTGGCCCGG TCGGCACCAAG TTGCGTGAGC GGAAAGATGG
CCGCTTCCCG GCCCTGCTGC AGGGAGCTCA AAATGGAGGA CGCGGCGCTC GGGAGAGCGG GCGGGTGAGT
CACCCACACA AAGGAAAAGG GCCTTCCCGT CCTCAGCCGT CGCTTCATGT GACTCCACGG AGTACCGGGC
GCCGTCCAGG CACCTCGATT AGTTCTCGAG CTTTGGAGT ACGTCGCTT TAGGTTGGGG GGAGGGGTTT
TATGCGATGG AGTTTCCCCA CACTGAGTGG GTGGAGACTG AAGTTAGGCC AGCTTGGCAG TTGATGTAAT
TCTCCTTGGG ATTTGCCCTT TTTGAGTTTG GATCTTGTTT CATTCTCAAG CCTCAGACAG TGGTTCAAAG
TTTTTTCTT CCATTTCAAG TGTCGTGAAT GGAGAGCGAC GAGAGCGGCC TGCCCGCCAT GGAGATCGAG
TGCCGCATCA CCGGCACCTT GAACGGCGTG GAGTTCGAGC TGGTGGGCGG CGGAGAGGGC ACCCCGAGC
AGGGCCGCAT GACCAACAAG ATGAAGAGCA CCAAAGGCGC CCTGACCTTC AGCCCTACC TGCTGAGCCA
CGTGATGGG TACGGTTCTT ACCACTTCGG CACCTACCCC AGCGGCTACG AGAACCCCTT CCTGCACGCC
ATCAACAACG GCGGCTACAC CAACACCCG ATCGAGAAGT ACGAGGACGG CGGCGTGCTG CACGTGAGCT
TCAGCTACCG CTACGAGGCC GGCCGCGTGA TCGGCGACTT CAAGGTGATG GGCACCGGCT TCCCCGAGGA
CAGCGTGATC TTCACCGACA AGATCATCCG CAGCAACGCC ACCGTGGAGC ACCTGCACCC CATGGCGGAT
AACGATCTGG ATGGCAGCTT CACCCGACCC TTCAGCCTGC GCGACGGCGG CTAATACAGC TCCGTGGTGG
ACAGCCACAT GCACTTCAAG AGCGCCATCC ACCCCAGCAT CCTGCAGAAC GGGGGCCCCA TGTTCCGCTT
CCGCCCGGTG GAGGAGGATC ACAGCAACAC CGAGCTGGGC ATCGTGAGT ACCAGCACGC CTTCAAGACC
CCGGATGCAG ATGCCGGTGA AGAAAGAGGA AGCGGAGCTA CTAAGTTCAG CTTGCTGAAG CAGGCTGGAG
ACGTGGAGGA GAACCTTGA CCTATGACCG AGTACAAGCC CACGGTGCGC CTCGCCACCC GCGACGACGT
CCCCAGGGCC GTACGCACCC TCGCCGCCGC GTTCGCCGAC TACCCGCA CGCGCCACAC CGTCGATCCG
GACCGCCACA TCGAGCGGGT CACCGAGCTG CAAGAACTCT TCCTCACGCG CGTCGGGCTC GACATCGGCA
AGGTGTGGT GCGGACGAC GCGCGCCGCG TGGCGGTCTG GACCACGCCG GAGAGCGTGC AAGCGGGGGC
GGTGTTCGCC GAGATCGGCC CGCGCATGGC CGAGTTGAGC GGTTCGCGG TGCGCCGCGA GCAACAGATG
GAAGGCCCTC TGCGCCGCA CCGGCCAAG GAGCCCGCTG GGTTCCTGGC CACCGTCGGC GTCTCGCCG
ACCACAGGG CAAGGTCTG GGCAGCGCCG TCGTGCTCC CGAGTGAGG GCGGCCGAGC GCGCCGGGT
GCCCGCTTC CTGGAGACCT CCGCGCCCG CAACCTCCCC TTCTACGAG GGCTCGGCTT CACCGTCACC
GCCGACGTC AGGTGCCGA AGGACCGCGC ACCTGGTGCA TGACCCGCAA GCCCGGTGCC TGAACTTGT
TTATTGCAGC TTATAATGGT TACAAATAA GCAATAGCAT CACAAATTTT ACAAATAAAG CATTTTTTTC
ACTGCATTCT AGTTGTGGT TGTCCAACT CATCAATGTA TCTTAATAA TTCGTATAAT GTATGCTATA CGAAGTTAT

```



RefSeq: [NM_008645](#)

UniProt ID: [P28665](#)

Summary: A proteinase activates the inhibitor by specific proteolysis in the bait region, which, by an unknown mechanism leads to reaction at the cysteinyl-glutamyl internal thiol ester site and to a conformational change, whereby the proteinase is trapped and/or covalently bound to the inhibitor. While in the tetrameric proteinase inhibitors steric inhibition is sufficiently strong, monomeric forms need a covalent linkage between the activated glutamyl residue of the original thiol ester and a terminal amino group of a lysine or another nucleophilic group on the proteinase, for inhibition to be effective.[UniProtKB/Swiss-Prot Function]

Product images:

