

Product datasheet for **KN513456**

Plekhg6 Mouse Gene Knockout Kit (CRISPR)

Product data:

Product Type:	Knockout Kits (CRISPR)
Format:	2 gRNA vectors, 1 linear donor
Donor DNA:	EF1a-GFP-P2A-Puro
Symbol:	Plekhg6
Locus ID:	213522



[View online »](#)

Components:
KN513456G1, Plekhg6 gRNA vector 1 in pCas-Guide CRISPR vector (GE100002)

KN513456G2, Plekhg6 gRNA vector 2 in pCas-Guide CRISPR vector (GE100002)

KN513456D, Linear donor DNA containing LoxP-EF1A-tGFP-P2A-Puro-LoxP:

The sequence below is cassette sequence only. The linear donor DNA also contains proprietary target sequence.

LoxP-EF1A-tGFP-P2A-Puro-LoxP (2739 bp)

```

ATAACTTCGT ATAATGTATG CTATACGAAG TTATCGTGAG GCTCCGGTGC CCGTCAGTGG GCAGAGCGCA
CATCGCCAC AGTCCCGAG AAGTTGGGG GAGGGGTGCG CAATTGAACC GGTGCCTAGA GAAGGTGGCG
CGGGGTAAAC TGGGAAAGTG ATGTCGTGTA CTGGCTCCGC CTTTTTCCCG AGGGTGGGGG AGAACCGTAT
ATAAGTCAG TAGTCGCCGT GAACGTTCTT TTTGCAACG GGTTCGCCG CAGAACACAG GTAAGTGCCG
TGTGTGGTTC CCGCGGGCCT GGCCTCTTTA CGGGTTATGG CCCTTGCGTG CCTTGAATTA CTTCCACCTG
GCTGCAGTAC GTGATTCTTG ATCCCGAGCT TCGGGTTGGA AGTGGGTGGG AGAGTTCGAG GCCTTGCGCT
TAAGGAGCCC CTTCGCCTCG TGCTTGAGTT GAGGCCTGGC CTGGGCGCTG GGGCCGCCG GTGCGAATCT
GGTGGCACCT TCGCGCCTGT CTCGCTGCTT TCGATAAGTC TCTAGCCATT TAAAATTTT GATGACCTGC
TGCAGCGCTT TTTTCTGGC AAGATAGTCT TGTAAATGCG GGCCAAGATC TGCACACTGG TATTTTCGTT
TTTGGGGCCG CGGGCGGCGA CGGGGCCCGT GCGTCCCAGC GCACATGTTC GGCAGGCGG GGCCTGCGAG
CGCGGCCACC GAGAATCGGA CGGGGGTAGT CTCAAGCTGG CCGGCCTGCT CTGGTGCCTG GCCTCGCGCC
GCCGTGTATC GCCCGGCCCT GGGCGGCAAG GCTGGCCCGG TCGGCACCAAG TTGCGTGAGC GGAAAGATGG
CCGCTTCCCG GCCCTGCTGC AGGGAGCTCA AAATGGAGGA CGCGGCGCTC GGGAGAGCGG GCGGGTGAGT
CACCCACACA AAGGAAAAGG GCCTTTCCGT CCTCAGCCGT CGCTTCATGT GACTCCACGG AGTACCGGGC
GCCGTCCAGG CACCTCGATT AGTTCTCGAG CTTTGGAGT ACCTCGTCTT TAGGTTGGGG GGAGGGGTTT
TATGCGATGG AGTTTCCCCA CACTGAGTGG GTGGAGACTG AAGTTAGGCC AGCTTGGCAG TTGATGTAAT
TCTCCTTGGG ATTTGCCCTT TTTGAGTTG GATCTTGGTT CATTCTCAAG CCTCAGACAG TGGTTCAAAG
TTTTTTCTT CCATTTCAAG TGTCGTGAAT GGAGAGCGAC GAGAGCGGCC TGCCCGCCAT GGAGATCGAG
TGCCGCATCA CCGGCACCTT GAACGGCGTG GAGTTCGAGC TGGTGGGCGG CGGAGAGGGC ACCCCGAGC
AGGGCCGCAT GACCAACAAG ATGAAGAGCA CCAAAGGCGC CCTGACCTTC AGCCCTACC TGCTGAGCCA
CGTGATGGG TACGGTTCTT ACCACTTCGG CACCTACCCC AGCGGCTACG AGAACCCCTT CCTGCACGCC
ATCAACAACG GCGGCTACAC CAACACCCG ATCGAGAAGT ACGAGGACGG CGGCGTGCTG CACGTGAGCT
TCAGCTACCG CTACGAGGCC GGCCGCGTGA TCGGCGACTT CAAGGTGATG GGCACCGGCT TCCCCGAGGA
CAGCGTGATC TTCACCGACA AGATCATCCG CAGCAACGCC ACCGTGGAGC ACCTGCACCC CATGGCGGAT
AACGATCTGG ATGGCAGCTT CACCCGCACC TTCAGCCTGC GCGACGGCGG CTAATACAGC TCCGTGGTGG
ACAGCCACAT GCACTTCAAG AGCGCCATCC ACCCCAGCAT CCTGCAGAAC GGGGGCCCCA TGTTGCCTT
CCGCCCGGTG GAGGAGGATC ACAGCAACAC CGAGCTGGGC ATCGTGAGT ACCAGCACGC CTTCAAGACC
CCGGATGCAG ATGCCGGTGA AGAAAGAGGA AGCGGAGCTA CTAAGTTTCA CCGTGTGAAG CAGGCTGGAG
ACGTGGAGGA GAACCTTGA CCTATGACCG AGTACAAGCC CACGGTGCGC CTCGCCACCC GCGACGACGT
CCCCAGGGCC GTACGCACCC TCGCCGCCGC GTTCGCCGAC TACCCGCA CGCGCCACAC CGTCGATCCG
GACCGCCACA TCGAGCGGGT CACCGAGCTG CAAGAACTCT TCCTCACGCG CGTCGGGCTC GACATCGGCA
AGGTGTGGT GCGGACGAC GCGCCCGCGG TGGCGGTCTG GACCACGCCG GAGAGCGTCTG AAGCGGGGGC
GGTGTTCGCC GAGATCGGCC CGCGCATGGC CGAGTTGAGC GGTTCGCGG TGCGCCGCA GCAACAGATG
GAAGGCCTCC TGGCGCCGCA CCGGCCAAG GAGCCCGCT GGTTCCTGGC CACCGTCGGC GTCTCGCCG
ACCACAGGG CAAGGTCTG GGCAGCGCG TCGTGCTCC CGAGTGAGG GCGGCCGAGC GCGCCGGGT
GCCCGCTTC CTGGAGACCT CCGCGCCCC CAACCTCCC TTCTACGAG GGCTCGGCTT CACCGTCACC
GCCGACGTC AGGTGCCGA AGGACCGCG ACCTGGTGCA TGACCCGCAA GCCCGGTGCC TGAACTTGT
TTATTGCAGC TTATAATGGT TACAAATAA GCAATAGCAT CACAAATTTT ACAAATAAAG CATTTTTTTC
ACTGCATTCT AGTTGTGGT TGTCCAACT CATCAATGTA TCTTAATAA TTCGTATAAT GTATGCTATA CGAAGTTAT

```



RefSeq: [NM_198604](#)

UniProt ID: [Q8R0J1](#)

Synonyms: BC026778; Gm155

Summary: Guanine nucleotide exchange factor activating the small GTPase RHOA, which, in turn, induces myosin filament formation. Also activates RHOG. Does not activate RAC1, or to a much lower extent than RHOA and RHOG. Part of a functional unit, involving PLEKHG6, MYH10 and RHOA, at the cleavage furrow to advance furrow ingression during cytokinesis. In epithelial cells, required for the formation of microvilli and membrane ruffles on the apical pole. Along with EZR, required for normal macropinocytosis (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:

