

## Product datasheet for **KN513868**

### Prkaa2 Mouse Gene Knockout Kit (CRISPR)

#### Product data:

Product Type:	Knockout Kits (CRISPR)
Format:	2 gRNA vectors, 1 linear donor
Donor DNA:	EF1a-GFP-P2A-Puro
Symbol:	Prkaa2
Locus ID:	108079



[View online »](#)

**Components:**
**KN513868G1**, Prkaa2 gRNA vector 1 in pCas-Guide CRISPR vector (GE100002)

**KN513868G2**, Prkaa2 gRNA vector 2 in pCas-Guide CRISPR vector (GE100002)

**KN513868D**, Linear donor DNA containing LoxP-EF1a-tGFP-P2A-Puro-LoxP:

The sequence below is cassette sequence only. The linear donor DNA also contains proprietary target sequence.

LoxP-EF1a-tGFP-P2A-Puro-LoxP (2739 bp)

```

ATAACTTCGT ATAATGTATG CTATACGAAG TTATCGTGAG GCTCCGGTGC CCGTCAGTGG GCAGAGCGCA
CATCGCCAC AGTCCCGAG AAGTTGGGG GAGGGTTCGG CAATTGAACC GGTGCCTAGA GAAGGTGGCG
CGGGTAAC TGGAAAGTG ATGTCGTGTA CTGGCTCCG CTTTTCCCG AGGGTGGGG AGAACCGTAT
ATAAGTCAG TAGTCGCCG GAACGTTCT TTTCCGAAC GGTTCGCCG CAGAACACAG GTAAGTGCCG
TGTGTGGTTC CCGCGGGCT GGCCTCTTA CGGGTTATG CCCTTGCGTG CCTTGAATTA CTTCCACCTG
GCTGCAGTAC GTGATTCTG ATCCCGAGCT TCGGGTTGGA AGTGGGTGG AGAGTTCGAG GCCTTGCGCT
TAAGGAGCCC CTTCGCCTG TGCTTGAGT GAGGCCTGG CTGGGCGTG GGGCCCGCG GTGCGAATCT
GGTGGCACCT TCGCGCCTG CTCGCTGCT TCGATAAGT TCTAGCCATT TAAAATTTT GATGACCTGC
TGCAGCGCT TTTTCTGGC AAGATAGTCT TGTAATGCG GGCCAAGATC TGCACACTGG TATTTTCGTT
TTTGGGGCCG CGGGCGGCA CGGGGCCCGT GCGTCCCAGC GCACATGTTC GGCGAGGCGG GGCCTGCGAG
CGCGGCCACC GAGAATCGGA CGGGGTAGT CCAAGCTGG CCGGCCTGCT CTGGTGCCTG GCCTCGCGCC
GCCGTGTATC GCCCGCCCT GGGCGGAAG GCTGGCCCG TCGGCACCAG TTGCGTGAGC GGAAAGATGG
CCGTTCCCG GCCCTGTGC AGGGAGCTCA AAATGGAGGA CGCGGCGCTC GGGAGAGCGG GCGGGTGAAGT
CACCCACACA AAGGAAAAG GCCTTCCGT CCTCAGCCG CGCTTCATGT GACTCCACGG AGTACCGGGC
GCCCTCCAG CACCTCGAT AGTTCTGAG CTTTGGAGT ACGTCGTCT TAGTTGGGG GGAGGGGTTT
TATGCGATGG AGTTCCCCA CACTGAGTG GTGAGACTG AAGTTAGGCC AGCTTGGCAG TTGATGTAAT
TCTCCTTGG AATTGCCCT TTTGAGTTG GATCTTGGT CATTCTCAAG CCTCAGACAG TGGTTCAAAG
TTTTTTCTT CCATTTCAAG TGTCGTGAAT GGAGAGCGAC GAGAGCGGCC TGCCCGCCAT GGAGATCGAG
TGCCGCATCA CCGGCACCCT GAACGGCGTG GAGTTCGAGC TGGTGGGCGG CGGAGAGGGC ACCCCCGAGC
AGGGCCGCAT GACCAACAAG ATGAAGAGCA CCAAAGGCGC CCTGACCTTC AGCCCTACC TGCTGAGCCA
CGTGATGGC TACGGCTTCT ACCACTTCG CACCTACCC AGCGGCTACG AGAACCCCTT CCTGCACGCC
ATCAACAACG GCGGCTACAC CAACACCCG ATCGAGAAGT ACGAGGACGG CGGCGTGCTG CACGTGAGCT
TCAGTACCG CTACGAGGC GGCCGCGTA TCGGCGACTT CAAGGTGATG GGCACCGCT TCCCGGAGGA
CAGCGTGATC TTCACCGACA AGATCATCCG CAGCAACGCC ACCGTGGAGC ACCTGCACCC CATGGCGAT
AACGATCTG ATGGCAGCT CACCCGACC TTCAGCCTG GCGACGGCG CTACTACAGC TCCGTGGTGG
ACAGCCACAT GCACTTCAAG AGCGCCATCC ACCCCAGCAT CCTGCAGAAC GGGGGCCCA TGTTCCCTT
CCGCCCGTG GAGGAGGATC ACAGCAACAC CGAGTGGGC ATCGTGGAGT ACCAGCACGC CTTCAAGACC
CCGGATGAG ATGCCGGTGA AGAAAGAGGA AGCGGAGCTA CTAACCTCAG CCTGCTGAAG CAGGCTGGAG
ACGTGGAGGA GAACCTGGA CCTATGACCG AGTACAAGC CACGGTGC GCCTGCCACC GCGACGACGT
CCCCAGGGC GTACGCACC TCGCCGCCG GTTCGCCGAC TACCCGCCA CGGCCACAC CGTCGATCCG
GACCGCCACA TCGAGCGGT CACCGAGCT CAAGAATCT TCCTCACGCG CGTCGGGCTC GACATCGCA
AGGTGTGGT CGCGGACGAC GGCGCCGCG TGGCGTCTG GACCACGCG GAGAGCGTCG AAGCGGGGGC
GGTGTTCGC GAGATCGCC CGCGCATGG CGAGTTGAGC GGTTCGGC TGGCCGCGCA GCAACAGATG
GAAGCCCTC TGCGCCGCA CCGGCCAAG GAGCCCGCT GTTCTCTGG CACCCTCGG GTCTCGCCG
ACCACCAGG CAAGGTCTG GGCAGCGCG TCGTGTCCC CGGAGTGGAG GCGGCCGAGC GCGCCGGGT
GCCCGCTT CTGGAGACT CCGCGCCCG CAACCTCCC TTCTACGAG GGCTCGGCT CACCGTCACC
GCCGACGTC AGGTGCCGA AGGACCGCG ACCTGGTGA TGACCCGCA GCCCGGTGCC TGAAACTTGT
TTATTGCAG TTATAATGG TACAAATAA GCAATAGCAT CACAAATTC ACAAATAAG CATTTTTTT
ACTGCATTCT AGTTGTGGT TGTCCAACT CATCAATGA TCTTAATAAC TTCGTATAAT GTATGTATA CGAAGTTAT
    
```



<b>Disclaimer:</b>	These products are manufactured and supplied by OriGene under license from ERS. The kit is designed based on the best knowledge of CRISPR technology. The system has been functionally validated for knocking-in the cassette downstream the native promoter. The efficiency of the knock-out varies due to the nature of the biology and the complexity of the experimental process.
<b>RefSeq:</b>	<a href="#">NM_178143</a> , <a href="#">NM_001356568</a>
<b>UniProt ID:</b>	<a href="#">Q8BRK8</a>
<b>Synonyms:</b>	231000811Rik; A830082D05; AMPKalpha2
<b>Summary:</b>	<p>Catalytic subunit of AMP-activated protein kinase (AMPK), an energy sensor protein kinase that plays a key role in regulating cellular energy metabolism. In response to reduction of intracellular ATP levels, AMPK activates energy-producing pathways and inhibits energy-consuming processes: inhibits protein, carbohydrate and lipid biosynthesis, as well as cell growth and proliferation. AMPK acts via direct phosphorylation of metabolic enzymes, and by longer-term effects via phosphorylation of transcription regulators. Also acts as a regulator of cellular polarity by remodeling the actin cytoskeleton; probably by indirectly activating myosin. Regulates lipid synthesis by phosphorylating and inactivating lipid metabolic enzymes such as ACACA, ACACB, GYS1, HMGCR and LIPE; regulates fatty acid and cholesterol synthesis by phosphorylating acetyl-CoA carboxylase (ACACA and ACACB) and hormone-sensitive lipase (LIPE) enzymes, respectively. Regulates insulin-signaling and glycolysis by phosphorylating IRS1, PFKFB2 and PFKFB3. Involved in insulin receptor/INSR internalization (By similarity). AMPK stimulates glucose uptake in muscle by increasing the translocation of the glucose transporter SLC2A4/GLUT4 to the plasma membrane, possibly by mediating phosphorylation of TBC1D4/AS160. Regulates transcription and chromatin structure by phosphorylating transcription regulators involved in energy metabolism such as CRTC2/TORC2, FOXO3, histone H2B, HDAC5, MEF2C, MLXIPL/ChREBP, EP300, HNF4A, p53/TP53, SREBF1, SREBF2 and PPARGC1A. Acts as a key regulator of glucose homeostasis in liver by phosphorylating CRTC2/TORC2, leading to CRTC2/TORC2 sequestration in the cytoplasm. In response to stress, phosphorylates 'Ser-36' of histone H2B (H2BS36ph), leading to promote transcription. Acts as a key regulator of cell growth and proliferation by phosphorylating TSC2, RPTOR and ATG1/ULK1: in response to nutrient limitation, negatively regulates the mTORC1 complex by phosphorylating RPTOR component of the mTORC1 complex and by phosphorylating and activating TSC2. In response to nutrient limitation, promotes autophagy by phosphorylating and activating ATG1/ULK1. In that process also activates WDR45. AMPK also acts as a regulator of circadian rhythm by mediating phosphorylation of CRY1, leading to destabilize it. May regulate the Wnt signaling pathway by phosphorylating CTNNB1, leading to stabilize it. Also phosphorylates CFTR, EEF2K, KLC1, NOS3 and SLC12A1. Plays an important role in the differential regulation of pro-autophagy (composed of PIK3C3, BECN1, PIK3R4 and UVRAG or ATG14) and non-autophagy (composed of PIK3C3, BECN1 and PIK3R4) complexes, in response to glucose starvation. Can inhibit the non-autophagy complex by phosphorylating PIK3C3 and can activate the pro-autophagy complex by phosphorylating BECN1 (PubMed:23332761). [UniProtKB/Swiss-Prot Function]</p>

Product images:

