

## Product datasheet for **LC400443**

### **FXYD6P3 (NM\_001099278) Human Over-expression Lysate**

#### **Product data:**

<b>Product Type:</b>	Over-expression Lysates
<b>Description:</b>	FXYD8 HEK293T cell transient overexpression lysate (as WB positive control)
<b>Species:</b>	Human
<b>Expression Host:</b>	HEK293T
<b>Expression cDNA Clone or AA Sequence:</b>	TrueORF Clone RC215461
<b>Tag:</b>	C-Myc/DDK
<b>Detection Antibodies:</b>	Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)
<b>ACCN:</b>	<a href="#">NM_001099278</a> , <a href="#">NP_001092748</a>
<b>Synonyms:</b>	FXYD6P3; FXYD domain-containing ion transport regulator 6, pseudogene 3; FXYD domain containing ion transport regulator 8; OTTHUMP00000023569
<b>Predicted MW:</b>	11 kDa
<b>Components:</b>	1 vial of 20 ug lyophilized gene specific transient over-expression cell lysate
<b>Storage:</b>	The lysate can be shipped at ambient temperature. Upon receiving, store the sample at -20°C. Lysate samples can be reconstituted with SDS Sample Buffer. Avoid repeated freeze-thaw cycles after reconstitution. Lysate samples are stable for 12 months from date of receipt when stored at -20°C.
<b>Preparation:</b>	HEK293T cells in 10-cm dishes were transiently transfected with <a href="#">MegaTran</a> Transfection Reagent (TT200002) and 5ug <a href="#">TrueORE</a> cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). To facilitate transportation and protein, the products are supplied as lyophilized proteins.
<b>RefSeq:</b>	<a href="#">NP_001092748</a>
<b>Locus ID:</b>	406875
<b>Cytogenetics:</b>	Xq13.2
<b>Protein Families:</b>	Transmembrane



[View online »](#)