

Product datasheet for **LC408018**


MAX (NM_145112) Human Over-expression Lysate

Product data:

Product Type:	Over-expression Lysates
Description:	MAX HEK293T cell transient overexpression lysate (as WB positive control)
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	TrueORF Clone RC206812
Tag:	C-Myc/DDK
Detection Antibodies:	Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)
ACCN:	NM_145112 , NP_660087
Synonyms:	bHLHd4
Predicted MW:	17.2 kDa
Components:	1 vial of 20 ug lyophilized gene specific transient over-expression cell lysate
Storage:	The lysate can be shipped at ambient temperature. Upon receiving, store the sample at -20°C. Lysate samples can be reconstituted with SDS Sample Buffer. Avoid repeated freeze-thaw cycles after reconstitution. Lysate samples are stable for 12 months from date of receipt when stored at -20°C.
Preparation:	HEK293T cells in 10-cm dishes were transiently transfected with MegaTran Transfection Reagent (TT200002) and 5ug TrueORE cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). To facilitate transportation and protein, the products are supplied as lyophilized proteins.
RefSeq:	NP_660087
Locus ID:	4149
Cytogenetics:	14q23.3
Protein Families:	Druggable Genome, Transcription Factors
Protein Pathways:	MAPK signaling pathway, Pathways in cancer, Small cell lung cancer



[View online »](#)

Product images:

DNA-binding activity of MAX was measured in OriGene over-expression lysate [LY408018] and a control lysate. Three microliters of each lysate was tested with a transcription factor binding assay utilizing MAX-specific DNA sequences. The high level of activity observed in the over-expression lysate compared to the control lysate demonstrates that the expressed MAX is biologically active in the lysate. Overexpression cell lysates are prepared from HEK293T cells transfected with [RC206812] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).