

Product datasheet for LC416187

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

UGT1A4 (NM 007120) Human Over-expression Lysate

Product data:

Product Type: Over-expression Lysates

Description: UGT1A4 HEK293T cell transient overexpression lysate (as WB positive control)

Species: Human HEK293T **Expression Host:**

Expression cDNA Clone

TrueORF Clone RC219801 or AA Sequence:

Tag:

C-Myc/DDK

Detection Antibodies: Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)

ACCN: NM 007120, NP 009051

GNT1; hUG-BR1; HUG-BR2; UDPGT; UDPGT 1-4; UGT-1A; UGT-1D; UGT1; UGT1-01; UGT1-04; Synonyms:

UGT1.1; UGT1.4; UGT1A; UGT1A1; UGT1A4S; UGT1D

Predicted MW: 60.03 kDa

1 vial of 20 ug lyophilized gene specific transient over-expression cell lysate Components:

The lysate can be shipped at ambient temperature. Upon receiving, store the sample at -Storage:

> 20°C. Lysate samples can be reconstituted with SDS Sample Buffer. Avoid repeated freezethaw cycles after reconstitution. Lysate samples are stable for 12 months from date of receipt

when stored at -20°C.

Preparation: HEK293T cells in 10-cm dishes were transiently transfected withMegaTran Transfection

> Reagent (TT200002) and 5ug TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). To facilitate transportation and protein, the

products are supplied as lyophilized proteins.

RefSeq: NP 009051

Locus ID: 54657 Cytogenetics: 2q37.1

Protein Families: Transmembrane






Protein Pathways:

Androgen and estrogen metabolism, Ascorbate and aldarate metabolism, Drug metabolism - cytochrome P450, Drug metabolism - other enzymes, Metabolic pathways, Metabolism of xenobiotics by cytochrome P450, Pentose and glucuronate interconversions, Porphyrin and chlorophyll metabolism, Retinol metabolism, Starch and sucrose metabolism

Product images:



Western blot validation of overexpression lysate (Cat# [LY416187]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with [RC219801] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).