

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for LC416620

GAS2L1 (NM_006478) Human Over-expression Lysate


Product data:

e: Over-expression	
	Lysates
GAS2L1 HEK293	cell transient overexpression lysate (as WB positive control)
Human	
Host: HEK293T	
:DNA Clone TrueORF Clone R ence:	C203825
C-Myc/DDK	
ntibodies: Clone OTI4C5, Ar	nti-DDK (FLAG) monoclonal antibody (TA50011-100)
<u>NM 006478, NP</u>	006469
GAR22	
W: 72.7 kDa	
s: 1 vial of 20 ug lyo	ophilized gene specific transient over-expression cell lysate
20°C. Lysate sam	e shipped at ambient temperature. Upon receiving, store the sample at - pples can be reconstituted with SDS Sample Buffer. Avoid repeated freeze- reconstitution. Lysate samples are stable for 12 months from date of receipt 20°C.
Reagent (TT2000 48hrs before col 150mM NaCl, 1% and 1mM Na3VC measured by BC	10-cm dishes were transiently transfected with <u>MegaTran</u> Transfection 02) and 5ug <u>TrueORF</u> cDNA plasmid. Transfected cells were cultured for lection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, o NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF 04), and then centrifuged to clarify the lysate. Protein concentration was A kit (Thermo Scientific Inc.). To facilitate transportation and protein, the oplied as lyophilized proteins.
<u>NP 006469</u>	
10634	
s: 22q12.2	
ence: C-Myc/DDK Clone OTI4C5, Ar <u>NM 006478, NP</u> GAR22 W: 72.7 kDa s: 1 vial of 20 ug lyo The lysate can be 20°C. Lysate sam thaw cycles after when stored at - : HEK293T cells in Reagent (TT2000 48hrs before col 150mM NaCl, 1% and 1mM Na3VC measured by BC products are sup <u>NP 006469</u> 10634	hti-DDK (FLAG) monoclonal antibody (TA50011-100) 006469 ophilized gene specific transient over-expression cell lysate e shipped at ambient temperature. Upon receiving, store the sample at - toples can be reconstituted with SDS Sample Buffer. Avoid repeated freeze- reconstitution. Lysate samples are stable for 12 months from date of rece 20°C. 10-cm dishes were transiently transfected with <u>MegaTran</u> Transfection 02) and Sug <u>TrueORF</u> cDNA plasmid. Transfected cells were cultured for lection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6 o NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMS 04), and then centrifuged to clarify the lysate. Protein concentration was A kit (Thermo Scientific Inc.). To facilitate transportation and protein, the



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Product images:



Western blot validation of overexpression lysate (Cat# [LY407671]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with [RC211855] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US