

Product datasheet for **LC419625**


EGR1 (NM_001964) Human Over-expression Lysate

Product data:

Product Type:	Over-expression Lysates
Description:	EGR1 HEK293T cell transient overexpression lysate (as WB positive control)
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	TrueORF Clone RC209956
Tag:	C-Myc/DDK
Detection Antibodies:	Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)
ACCN:	<u>NM_001964</u> , <u>NP_001955</u>
Synonyms:	AT225; G0S30; KROX-24; NGFI-A; TIS8; ZIF-268; ZNF225
Predicted MW:	57.3 kDa
Components:	1 vial of 20 ug lyophilized gene specific transient over-expression cell lysate
Storage:	The lysate can be shipped at ambient temperature. Upon receiving, store the sample at -20°C. Lysate samples can be reconstituted with SDS Sample Buffer. Avoid repeated freeze-thaw cycles after reconstitution. Lysate samples are stable for 12 months from date of receipt when stored at -20°C.
Preparation:	HEK293T cells in 10-cm dishes were transiently transfected with <u>MegaTran</u> Transfection Reagent (TT200002) and 5ug <u>TrueORE</u> cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). To facilitate transportation and protein, the products are supplied as lyophilized proteins.
RefSeq:	<u>NP_001955</u>
Locus ID:	1958
Cytogenetics:	5q31.2
Protein Families:	Druggable Genome
Protein Pathways:	Prion diseases



[View online »](#)

Product images:

DNA-binding activity of EGR1 was measured in OriGene over-expression lysate [LY419625] and a control lysate. Three microliters of each lysate was tested with a transcription factor binding assay utilizing EGR1-specific DNA sequences. The high level of activity observed in the over-expression lysate compared to the control lysate demonstrates that the expressed EGR1 is biologically active in the lysate. Overexpression cell lysates are prepared from HEK293T cells transfected with [RC209956] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).