

## **Product datasheet for LC422309**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## GGT1 (NM\_001032365) Human Over-expression Lysate

**Product data:** 

**Product Type:** Over-expression Lysates

**Description:** GGT1 HEK293T cell transient overexpression lysate (as WB positive control)

Species: Human
Expression Host: HEK293T

**Expression cDNA Clone** 

Expression CDNA Cione

TrueORF Clone RC223689

or AA Sequence:

Tag: C-Myc/DDK

**Detection Antibodies:** Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)

ACCN: <u>NM 001032365</u>, <u>NP 001027537</u>

**Synonyms:** CD224; D22S672; D22S732; GGT; GGT 1; GTG

**Predicted MW:** 61.2 kDa

**Components:** 1 vial of 20 ug lyophilized gene specific transient over-expression cell lysate

Storage: The lysate can be shipped at ambient temperature. Upon receiving, store the sample at -

20°C. Lysate samples can be reconstituted with SDS Sample Buffer. Avoid repeated freeze-thaw cycles after reconstitution. Lysate samples are stable for 12 months from date of receipt

when stored at -20°C.

**Preparation:** HEK293T cells in 10-cm dishes were transiently transfected with MegaTran Transfection

Reagent (TT200002) and 5ug <u>TrueORF</u> cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). To facilitate transportation and protein, the

products are supplied as lyophilized proteins.

**RefSeq:** NP 001027537

**Locus ID:** 2678

Cytogenetics: 22q11.23

**Protein Families:** Protease, Transmembrane







**Protein Pathways:** 

Arachidonic acid metabolism, Cyanoamino acid metabolism, Glutathione metabolism, Metabolic pathways, Selenoamino acid metabolism, Taurine and hypotaurine metabolism