

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for LC429589

## RAC1 (NM\_018890) Human Over-expression Lysate

## **Product data:**

| Product Type:                            | Over-expression Lysates  |
|--|--|
| Description:                             | RAC1 HEK293T cell transient overexpression lysate (as WB positive control)   |
| Species:                                 | Human  |
| Expression Host:                         | HEK293T  |
| Expression cDNA Clone<br>or AA Sequence: | TrueORF Clone RC224262   |
| Tag:                                     | C-Myc/DDK  |
| Detection Antibodies:                    | Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)  |
| ACCN:                                    | <u>NM 018890, NP 061485</u>  |
| Synonyms:                                | MIG5; MRD48; p21-Rac1; Rac-1; TC-25  |
| Predicted MW:                            | 23.3 kDa   |
| Components:                              | 1 vial of 20 ug lyophilized gene specific transient over-expression cell lysate  |
| Storage:                                 | The lysate can be shipped at ambient temperature. Upon receiving, store the sample at -<br>20°C. Lysate samples can be reconstituted with SDS Sample Buffer. Avoid repeated freeze-<br>thaw cycles after reconstitution. Lysate samples are stable for 12 months from date of receipt<br>when stored at -20°C.   |
| Preparation:                             | HEK293T cells in 10-cm dishes were transiently transfected with <u>MegaTran</u> Transfection<br>Reagent (TT200002) and 5ug <u>TrueORF</u> cDNA plasmid. Transfected cells were cultured for<br>48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6,<br>150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF<br>and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was<br>measured by BCA kit (Thermo Scientific Inc.). To facilitate transportation and protein, the<br>products are supplied as lyophilized proteins. |
| RefSeq:                                  | <u>NP 061485</u>   |
| Locus ID:                                | 5879   |
| Cytogenetics:                            | 7p22.1   |
| Protein Families:                        | Druggable Genome   |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US Protein Pathways:Adherens junction, Amyotrophic lateral sclerosis (ALS), Axon guidance, B cell receptor<br/>signaling pathway, Chemokine signaling pathway, Colorectal cancer, Epithelial cell signaling in<br/>Helicobacter pylori infection, Fc epsilon RI signaling pathway, Fc gamma R-mediated<br/>phagocytosis, Focal adhesion, Leukocyte transendothelial migration, MAPK signaling pathway,<br/>Natural killer cell mediated cytotoxicity, Neurotrophin signaling pathway, Pancreatic cancer,<br/>Pathways in cancer, Regulation of actin cytoskeleton, Renal cell carcinoma, Toll-like receptor<br/>signaling pathway, VEGF signaling pathway, Viral myocarditis, Wnt signaling pathway

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US