

Product datasheet for LC430462

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Phosphatidic acid phosphatase type 2B (PLPP3) (NM_177414) Human Over-expression Lysate

Product data:

Product Type: Over-expression Lysates

Description: PPAP2B HEK293T cell transient overexpression lysate (as WB positive control)

Species: Human
Expression Host: HEK293T

Expression cDNA Clone or AA Sequence:

TrueORF Clone RC224959

Tag:

C-Myc/DDK

Detection Antibodies: Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)

ACCN: <u>NM 177414, NP 803133</u>

Synonyms: Dri42; LPP3; MGC15306; PAP-2b; PAP2-beta; VCIP

Predicted MW: 34.9 kDa

Components: 1 vial of 20 ug lyophilized gene specific transient over-expression cell lysate

Storage: The lysate can be shipped at ambient temperature. Upon receiving, store the sample at -

20°C. Lysate samples can be reconstituted with SDS Sample Buffer. Avoid repeated freeze-thaw cycles after reconstitution. Lysate samples are stable for 12 months from date of receipt

when stored at -20°C.

Preparation: HEK293T cells in 10-cm dishes were transiently transfected with MegaTran Transfection

Reagent (TT200002) and 5ug <u>TrueORF</u> cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). To facilitate transportation and protein, the

products are supplied as lyophilized proteins.

RefSeq: NP 803133

Locus ID: 8613 Cytogenetics: 1p32.2

Protein Families: Druggable Genome, Transmembrane





Phosphatidic acid phosphatase type 2B (PLPP3) (NM_177414) Human Over-expression Lysate – LC430462

Protein Pathways:

Ether lipid metabolism, Fc gamma R-mediated phagocytosis, Glycerolipid metabolism, Glycerophospholipid metabolism, Metabolic pathways, Sphingolipid metabolism