

Product datasheet for LY400481

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

BMP2 (NM_001200) Human Over-expression Lysate

Product data:

Product Type: Over-expression Lysates

Description: Transient overexpression lysate of bone morphogenetic protein 2 (BMP2)

Species: Human
Expression Host: HEK293T

Expression cDNA Clone

TrueORF Clone RC214586

or AA Sequence:

Tag: C-Myc/DDK

Detection Antibodies: Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)

ACCN: <u>NM 001200, NP 001191</u>

Synonyms: BDA2; BMP2A; SSFSC; SSFSC1

Predicted MW: 44.63 kDa

Components: 1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer

1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer

1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%

Bromophenol blue, 100mM DTT)

Storage: The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after

dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at -

80°C.

Preparation: HEK293T cells in 10-cm dishes were transiently transfected with MegaTran Transfection

Reagent (TT200002) and 5ug <u>TrueORF</u> cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was

measured by BCA kit (Thermo Scientific Inc.).

RefSeq: NP 001191

Locus ID: 650

Cytogenetics: 20p12.3





Protein Families:


WB

Protein Pathways: Acute myeloid leukemia, Basal cell carcinoma, Cytokine-cytokine receptor interaction,

Endocytosis, Hedgehog signaling pathway, Hematopoietic cell lineage, Melanogenesis,

Pathways in cancer, TGF-beta signaling pathway

Product images:



Western blot validation of overexpression lysate (Cat# LY400481) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with [RC214586] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).