

Product datasheet for LY400630

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

RHOA (NM 001664) Human Over-expression Lysate

Product data:

Product Type: Over-expression Lysates

Description: Transient overexpression lysate of ras homolog gene family, member A (RHOA)

Species: Human HEK293T **Expression Host:**

Expression cDNA Clone

or AA Sequence:

TrueORF Clone RC203303

Tag: C-Myc/DDK

Detection Antibodies: Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)

ACCN: NM 001664, NP 001655

ARH12; ARHA; EDFAOB; RHO12; RHOH12 Synonyms:

Predicted MW: 21.8 kDa

1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer Components:

1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer

1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%

Bromophenol blue, 100mM DTT)

The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after Storage:

> dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at -

80°C.

Preparation: HEK293T cells in 10-cm dishes were transiently transfected with MegaTran Transfection

> Reagent (TT200002) and 5ug TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). Cell lysates were aliquoted and stored at -20°C

before shipping.

RefSeq: NP 001655

Locus ID: 387





Cytogenetics: 3p21.31


Protein Families: Druggable Genome

Protein Pathways: Adherens junction, Axon guidance, Chemokine signaling pathway, Focal adhesion, Leukocyte

transendothelial migration, Neurotrophin signaling pathway, Pathogenic Escherichia coli infection, Pathways in cancer, Regulation of actin cytoskeleton, T cell receptor signaling pathway, TGF-beta signaling pathway, Tight junction, Vascular smooth muscle contraction,

Wnt signaling pathway

Product images:



Western blot validation of overexpression lysate (Cat# LY400630) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with [RC203303] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).