

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for LY402014

### PSKH1 (NM\_006742) Human Over-expression Lysate

### **Product data:**


Description:Transient overexpression lysate of protein serine kinase H1 (PSKH1)Species:HumanExpression Host:HEK293TExpression cDNA Cloop or AA Sequence:TrueORF Clone RC208968Tag:O-My/DDKDetection Antibodies:Cone OT14C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)ACCN:M. 006742, NP 006733Predicted MW:48 kDaComponents:Viai of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 viai of 100 µg gene specific transient over-expression cell syste in RIPA buffer 1 viai of 100 µg gene specific transient over-expression cell syste in RIPA buffer 1 viai of 100 µg sene specific transient over-expression cell syste in RIPA buffer 1 viai of 100 µg sene specific transient over-expression cell syste in RIPA buffer 1 viai of 100 µg sene specific transient over-expression cell syste in SIPA buffer 1 viai of 100 µg whole HEK293T cell syste in RIPA buffer 1 viai of 100 µg whole HEK293T cell syste in subject with dy to 2500 and specific transient over-expression cell syste in SIPA buffer 1 viai of 2500 µz SDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTTStorage:The lysate is shipped with dy ice. Upon receiving, store the sample at-80°C, Also after and trito, the protein sample should be aliquoted and store at-80°C forlong term storage, Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer (4% SDS, Also After Somple Assert CT20002) and Sug TrueORF CDNA plasmid. Transfetced cells were cultured of stores coll in modified RIPA buffer (25mM Tris-HCl pH7.2) SomM NAC1, MNP4.0, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMS Sigma, 1mM NAS4/A) and then centrifuged to clarify the lysa	Product Type:	Over-expression Lysates
Expression Host:HEK293TExpression cDNA Clone or AA Sequence:TrueORF Clone RC208968Tag:C-Myc/DDKDetection Antibodies:Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)ACCN:NM 006742, NP 006733Predicted MW:48 kDaComponents:1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell spate in RIPA buffer 2 bow or phenol blue, 100mM DTT)Storage:The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provide. Lysate samples are stable for 12 months from the date of receipt when stored at 80°C.Preparation:HEK293T cells in 10-cm dishes were transiently transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCI pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMS- and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). Cell lysates were aliquoted and stored at -20°C before shipping.RefSeq:NP 006733Locus ID:5681	Description:	Transient overexpression lysate of protein serine kinase H1 (PSKH1)
Expression cDNA CloneTrueORF Clone RC208968Tag:C-Myc/DDKDetection Antibodies:Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)ACCN:NM 006742, NP 006733Predicted MW:48 kDaComponents:1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)Storage:The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer orvided. Lysate samples are stable for 12 months from the date of receipt when stored at 80°C.Preparation:HEK293T cells in 10-cm dishes were transiently transfected cells were cultured for Alsn's before collection. The cells were lysade in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). Cell lysates were aliquoted ant stored at -20°C before shipping.RefSeq:NP.006733Locus ID:5681	Species:	Human
or AA Sequence:Tag:C-Myc/DDKDetection Antibodies:Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)ACCN:NM 006742, NP 006733Predicted MW:48 kDaComponents:1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)Storage:The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C Also after dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer rovided. Lysate samples are stable for 12 months from the date of receipt when stored at s0°C.Preparation:HEK293T cells in 10-cm dishes were transiently transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.4 SomM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). Cell lysates were aliquoted and stored at -20°C before shipping.RefSeq:NP 006733Locus ID:5681	Expression Host:	HEK293T
Detection Antibodies:Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)ACCN:NM 006742, NP 006733Predicted MW:48 kDaComponents:1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)Storage:The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at -80°C.Preparation:HEK293T cells in 10-cm dishes were transiently transfected withMegaTran Transfection Reagent (TT200002) and Sug TrueORE cDNA plasmid. Transfected cells were cultured for A8hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). Cell lysates were aliquoted and stored at -20°C before shipping.RefSeq:NP 006733Locus ID:5681	•	TrueORF Clone RC208968
ACCN:NM 006742, NP 006733Predicted MW:48 kDaComponents:1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)Storage:The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at - 80°C.Preparation:HEK293T cells in 10-cm dishes were transiently transfected withMegaTran Transfection Reagent (TT20002) and Sug TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). Cell lysates were aliquoted and stored at -20°C before shipping.RefSeq:NP 006733Locus ID:5681	Tag:	C-Myc/DDK
Predicted MW:48 kDaComponents:1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)Storage:The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at - 80°C.Preparation:HEK293T cells in 10-cm dishes were transiently transfected withMegaTran Transfection Reagent (TT200002) and Sug TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). Cell lysates were aliquoted and stored at -20°C before shipping.RefSeq:NP 006733Locus ID:5681	<b>Detection Antibodies:</b>	Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)
Components:1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)Storage:The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at -80°C.Preparation:HEK293T cells in 10-cm dishes were transiently transfected withMegaTran Transfection Reagent (TT200002) and 5ug TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). Cell lysates were aliquoted and stored at -20°C before shipping.RefSeq:NP 006733 E061	ACCN:	<u>NM 006742, NP 006733</u>
1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)Storage:The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at - 80°C.Preparation:HEK293T cells in 10-cm dishes were transiently transfected withMegaTran Transfection Reagent (TT200002) and 5ug TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). Cell lysates were aliquoted and stored at -20°C before shipping.RefSeq:NP 006733Locus ID:5681	Predicted MW:	48 kDa
<ul> <li>dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at -80°C.</li> <li>Preparation: HEK293T cells in 10-cm dishes were transiently transfected withMegaTran Transfection Reagent (TT200002) and 5ug TrueORE cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). Cell lysates were aliquoted and stored at -20°C before shipping.</li> <li>RefSeq: NP 006733</li> <li>Locus ID: 5681</li> </ul>	Components:	1 vial of 100 μg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%
Reagent (TT200002) and 5ug TrueORE cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). Cell lysates were aliquoted and stored at -20°C before shipping.RefSeq:NP 006733Locus ID:5681	Storage:	dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at -
Locus ID: 5681	Preparation:	Reagent (TT200002) and 5ug <u>TrueORF</u> cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). Cell lysates were aliquoted and stored at -20°C
	RefSeq:	<u>NP 006733</u>
Cytogenetics: 16q22.1	Locus ID:	5681
	Cytogenetics:	16q22.1



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US Protein Families:

Druggable Genome, Protein Kinase

## Product images:



Western blot validation of overexpression lysate (Cat# LY402014) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with [RC208968] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US