

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for LY405455

### ZNF677 (NM\_182609) Human Over-expression Lysate

#### **Product data:**


Description:Iransient overexpression lysate of zinc finger protein 677 (ZNF677)Species:HumanExpression Host:HEK293TComponents:Onyc/DDKDetection Antibodies:Components:Vial of 100 µg gene specific transient over-expression cell ysate in RIPA buffer i vial of 100 µg sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycenot, 00000)	Product Type:	Over-expression Lysates
Expression Host:HEK293TExpression cDNA Clone or AA Sequence:TrueORF Clone RC207997Tag:C-Myc/DDKDetection Antibodies:Cone OT14C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)ACCN:NM 182609, NP 872415Predicted MW:68 kDaComponents:1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%	Description:	Transient overexpression lysate of zinc finger protein 677 (ZNF677)
Expression cDNA Clone or AA Sequence:TrueORF Clone RC207997Tag:C-Myc/DDKDetection Antibodies:Cone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)ACCN:NM 182609, NP 872415Predicted MW:68 kDaComponents:1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%	Species:	Human
or AA Sequence:Tag:C-Myc/DDKDetection Antibodies:Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)ACCN:NM 182609, NP 872415Predicted MW:68 kDaComponents:1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%	Expression Host:	HEK293T
Detection Antibodies:Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)ACCN:NM 182609, NP 872415Predicted MW:68 kDaComponents:1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%	•	TrueORF Clone RC207997
ACCN:NM 182609, NP 872415Predicted MW:68 kDaComponents:1 vial of 100 μg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 μg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%)	Tag:	C-Myc/DDK
Predicted MW: 68 kDa   Components: 1 vial of 100 μg gene specific transient over-expression cell lysate in RIPA buffer   1 vial of 100 μg whole HEK293T cell lysate in RIPA buffer   1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%)	<b>Detection Antibodies:</b>	Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)
Components:1 vial of 100 μg gene specific transient over-expression cell lysate in RIPA buffer1 vial of 100 μg whole HEK293T cell lysate in RIPA buffer1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%)	ACCN:	<u>NM 182609, NP 872415</u>
1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%	Predicted MW:	68 kDa
Bromophenol blue, 100mM DTT)	Components:	1 vial of 100 μg whole HEK293T cell lysate in RIPA buffer
Storage:The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at - 	Storage:	dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at -
Preparation:HEK293T cells in 10-cm dishes were transiently transfected with MegaTran Transfection Reagent (TT200002) and 5ug TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). Cell lysates were aliquoted and stored at -20°C 	Preparation:	Reagent (TT200002) and 5ug <u>TrueORF</u> cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). Cell lysates were aliquoted and stored at -20°C
<b>RefSeq:</b> <u>NP 872415</u>	RefSeq:	<u>NP 872415</u>
Locus ID: 342926	Locus ID:	342926
<b>Cytogenetics:</b> 19q13.42	Cytogenetics:	19q13.42



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US Protein Families: Trai

Transcription Factors

## **Product images:**



Western blot validation of overexpression lysate (Cat# LY405455) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with [RC207997] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US