

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for LY405959

LCE3E (NM_178435) Human Over-expression Lysate

Product data:

Description:Transient overexpression lysate of late cornified envelope 3E (LCE3E)Species:HumanExpression Host:HEK293TExpression cDNA Clone or AA Sequence:TrueORF Clone RC213561Tag:C-Myc/DDKDetection Antibodies:Cone OT14C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)ACCN:M1 178435, NP 848522Synonyms:LEP17Predicted MW:9.5 kDaComponents:Nial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 25001 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%) Bromphene Juice Spinol blue, 100mM DTTStorage:The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after
Fxpression Host:HEK293TExpression cDNA Clooe or AA Sequence:TrueORF Clone RC213561Tag:C-Myc/DDKDetection Antibodies:Cone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)ACCN:NM 178435, NP 848522Synonyms:LEP17Predicted MW:9.5 kDaComponents:1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%) Bromphenol blue, 100mM DTT)
Expression cDNA CloneTrueORF Clone RC213561Tag:C-Myc/DDKTag:C-Myc/DDKDetection Antibodies:Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)ACCN:MM 178435, NP 848522Synonyms:LEP17Predicted MW:9.5 kDaComponents:1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%) Bromophenol blue, 100mM DTT
or AA Sequence:Tag:C-Myc/DDKDetection Antibodies:Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)ACCN:NM 178435, NP 848522Synonyms:LEP17Predicted MW:9.5 kDaComponents:1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%)
Detection Antibodies:Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)ACCN:NM 178435, NP 848522Synonyms:LEP17Predicted MW:9.5 kDaComponents:1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)
ACCN:NM 178435, NP 848522Synonyms:LEP17Predicted MW:9.5 kDaComponents:1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)
Synonyms:LEP17Predicted MW:9.5 kDaComponents:1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)
Predicted MW:9.5 kDaComponents:1 vial of 100 μg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 μg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)
Components:1 vial of 100 μg gene specific transient over-expression cell lysate in RIPA buffer1 vial of 100 μg whole HEK293T cell lysate in RIPA buffer1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%Bromophenol blue, 100mM DTT)
1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)
Storage: The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after
dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at - 80°C.
Preparation:HEK293T cells in 10-cm dishes were transiently transfected with MegaTran Transfection Reagent (TT200002) and 5ug TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). Cell lysates were aliquoted and stored at -20°C before shipping.
RefSeq: <u>NP 848522</u>
Locus ID: 353145



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US Cytogenetics: 1q21.3

Product images:

158-	
106 - 79 -	
48 -	
35 - 23 -	

Western blot validation of overexpression lysate (Cat# LY405959) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with [RC213561] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US