

## Product datasheet for LY406700

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## IL4I1 (NM 172374) Human Over-expression Lysate

**Product data:** 

**Product Type:** Over-expression Lysates

**Description:** Transient overexpression lysate of interleukin 4 induced 1 (IL4I1), transcript variant 2

Species: Human HEK293T **Expression Host:** 

**Expression cDNA Clone** 

or AA Sequence:

TrueORF Clone RC221972

Tag: C-Myc/DDK

**Detection Antibodies:** Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)

ACCN: NM 172374, NP 758962 FIG1; hIL4I1; LAAO; LAO Synonyms:

**Predicted MW:** 65.4 kDa

1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer Components:

1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer

1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%

Bromophenol blue, 100mM DTT)

The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after Storage:

> dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at -

80°C.

Preparation: HEK293T cells in 10-cm dishes were transiently transfected with MegaTran Transfection

> Reagent (TT200002) and 5ug TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was

measured by BCA kit (Thermo Scientific Inc.).

NP 758962 RefSeq:

Locus ID: 259307

Cytogenetics: 19q13.33






**Protein Families:** 

WB

**Protein Pathways:** 

Alanine, aspartate and glutamate metabolism, Cysteine and methionine metabolism, Metabolic pathways, Phenylalanine, tyrosine and tryptophan biosynthesis, Phenylalanine metabolism, Tryptophan metabolism, Tyrosine metabolism, Valine, leucine and isoleucine degradation

## **Product images:**



Western blot validation of overexpression lysate (Cat# LY406700) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with [RC221972] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).