

## Product datasheet for LY415713

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## TBK1 (NM 013254) Human Over-expression Lysate

**Product data:** 

**Product Type:** Over-expression Lysates

**Description:** Transient overexpression lysate of TANK-binding kinase 1 (TBK1)

TrueORF Clone RC205238

Species: Human HEK293T **Expression Host:** 

**Expression cDNA Clone** 

or AA Sequence:

Tag: C-Myc/DDK

**Detection Antibodies:** Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)

ACCN: NM 013254, NP 037386 FTDALS4; IIAE8; NAK; T2K Synonyms:

**Predicted MW:** 83.6 kDa

1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer Components:

1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer

1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%

Bromophenol blue, 100mM DTT)

The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after Storage:

> dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at -

80°C.

Preparation: HEK293T cells in 10-cm dishes were transiently transfected with MegaTran Transfection

> Reagent (TT200002) and 5ug TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was

measured by BCA kit (Thermo Scientific Inc.).

NP 037386 RefSeq:

Locus ID: 29110 Cytogenetics: 12q14.2





## TBK1 (NM\_013254) Human Over-expression Lysate - LY415713

Protein Families: WB

Protein Pathways: Cytosolic DNA-sensing pathway, RIG-I-like receptor signaling pathway, Toll-like receptor

signaling pathway