

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for LY418569

#### Aurora A (AURKA) (NM\_003600) Human Over-expression Lysate

### **Product data:**


Description:Transient overexpression lysate of aurora kinase A (AURKA), transcript variant 2Species:HumanExpression Host:HEK293TExpression cDNA Clone or AA Sequence:TrueORF Clone RC216460Tag:C-Myc/DDKDetection Antibodies:Cone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)ACCN:NM 003600, NP 003591Synonyms:AiK; ARK1; AURA; BTAK; PPP1R47; STK6; STK7; STK15Predicted MW:45.6 kDaComponents:1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 2 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%) Borney Buffer (14% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%) Borney Buffer (14% SDS, 125mM Tris-HCl pH6.8, 10% Clycerol, 0.002%) Borney Buffer (14% SDS, 125mM Tris-HCl pH6.8, 10% Clycerol, 0.002%) Borney Buffer (14% SDS, 125mM Tris-HCl pH6.8, 10% Clycerol, 0.002%) Borney Buffer (14% SDS, 125mM Tris-HCl pH6.8, 10% Clycerol, 0.002%) Borney Buffer (14% SDS, 125mM Tris-HCl pH6.8, 10% Clycerol, 0.002%) Borney Buffer (14% SDS, 125mM Tris-HCl pH6.8, 10% Clycerol, 0.002%) Borney Buffer (14% SDS, 125mM Tris-HCl pH6.8, 10% Clycerol, 0.002%) Borney Buffer (14% SDS, 125mM Tris-HCl pH6.8, 10% Clycerol, 0.002%)
Fxpression Host:HEK293TExpression cDNA Clone or AA Sequence:TrueORF Clone RC216460Tag:C-Myc/DDKDetection Antibodies:Cone OT14C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)ACCN:NM 003600, NP 003591Synonyms:AIK; ARK1; AURA; BTAK; PPP1R47; STK6; STK7; STK15Predicted MW:45.6 kDaComponents:1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)Storage:The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after
Expression cDNA CloneTrueORF Clone RC216460or AA Sequence:C-Myc/DDKTag:C-Myc/DDKDetection Antibodies:Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)ACCN:NM 003600, NP 003591Synonyms:AIK; ARK1; AURA; BTAK; PPP1R47; STK6; STK7; STK15Predicted MW:45.6 kDaComponents:1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%) Bromophenol blue, 100mM DTT)Storage:The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after
or AA Sequence:Tag:C-Myc/DDKDetection Antibodies:Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)ACCN:NM 003600, NP 003591Synonyms:AIK; ARK1; AURA; BTAK; PPP1R47; STK6; STK7; STK15Predicted MW:45.6 kDaComponents:1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%) Bromophenol blue, 100mM DTT)Storage:The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after
Detection Antibodies:Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)ACCN:NM 003600, NP 003591Synonyms:AIK; ARK1; AURA; BTAK; PPP1R47; STK6; STK7; STK15Predicted MW:45.6 kDaComponents:1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)Storage:The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after
<ul> <li>ACCN: NM 003600, NP 003591</li> <li>Synonyms: AIK; ARK1; AURA; BTAK; PPP1R47; STK6; STK7; STK15</li> <li>Predicted MW: 45.6 kDa</li> <li>Components: 1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)</li> <li>Storage: The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after</li> </ul>
Synonyms:AIK; ARK1; AURA; BTAK; PPP1R47; STK6; STK7; STK15Predicted MW:45.6 kDaComponents:1 vial of 100 μg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 μg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)Storage:The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after
Predicted MW:45.6 kDaComponents:1 vial of 100 μg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 μg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)Storage:The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after
Components:1 vial of 100 μg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 μg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)Storage:The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after
1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%Bromophenol blue, 100mM DTT)Storage:The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after
dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at - 80°C.
Preparation:HEK293T cells in 10-cm dishes were transiently transfected with MegaTran Transfection Reagent (TT200002) and 5ug TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). Cell lysates were aliquoted and stored at -20°C before shipping.
<b>RefSeq:</b> <u>NP 003591</u>
Locus ID: 6790



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Aurora A (AURKA) (NM_003600) Human Over-expression Lysate – LY418569
Cytogenetics:	20q13.2
Protein Families: Protein Pathways	Druggable Genome, Protein Kinase, Stem cell - Pluripotency Oocyte meiosis

## **Product images:**



Western blot validation of overexpression lysate (Cat# LY418569) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with [RC216460] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US