

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for LY419328

## MSH3 (NM\_002439) Human Over-expression Lysate


## **Product data:**

Description:Transient overexpression lysate of mutS homolog 3 (E. coli) (MSH3)Species:HumanExpression Host:HEK293TExpression cDNA Clone or AA Sequence:TrueORF Clone RC212391Tag:C-Myc/DDKDetection Antibodies:Cone OT14C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)ACCN:MM 002439, NP 002430Synonyms:DUP; FAP4; MRP1Predicted MW:127.2 kDaComponents:1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCI pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)Storage:The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer violed. Lysate samples are stable for 12 months from the date of receipt when stored at a0°C.Preparation:HEK293T cells in 10-cm dishes were transiently transfected withMegaTran Transfection Reagent (TT200002) and Sug TrueORF cDNA plasmid. Transfected cells were cultured for
Expression Host:HEK293TExpression cDNA Cloop or AA Sequence:TrueORF Clone RC212391Tag:C-Myc/DDKDetection Antibodies:Clone OT14C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)ACCN:NM 002439, NP 002430Synonyms:DUP; FAP4; MRP1Predicted MW:127.2 kDaComponents:1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)Storage:The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer 12 months from the date of receipt when stored at -80°C.Preparation:HEK293T cells in 10-cm dishes were transiently transfected withMegaTran Transfection
Expression cDNA CloneTrueORF Clone RC212391Tag:C-Myc/DDKDetection Antibodies:Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)ACCN:NM 002439, NP 002430Synonyms:DUP; FAP4; MRP1Predicted MW:127.2 kDaComponents:1 vial of 100 µg gne specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)Storage:The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer avoid. Lysate samples are stable for 12 months from the date of receipt when stored at -80°C.Preparation:HEK293T cells in 10-cm dishes were transiently transfected withMegaTran Transfection
or AA Sequence:Tag:C-Myc/DDKDetection Antibodies:Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)ACCN:NM 002439, NP 002430Synonyms:DUP; FAP4; MRP1Predicted MW:127.2 kDaComponents:1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%) Bromophenol blue, 100mM DTT)Storage:The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at -80°C.Preparation:HEK293T cells in 10-cm dishes were transiently transfected withMegaTran Transfection
Detection Antibodies:Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)ACCN:NM 002439, NP 002430Synonyms:DUP; FAP4; MRP1Predicted MW:127.2 kDaComponents:1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)Storage:The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at - 80°C.Preparation:HEK293T cells in 10-cm dishes were transiently transfected withMegaTran Transfection
ACCN:NM 002439, NP 002430Synonyms:DUP; FAP4; MRP1Predicted MW:127.2 kDaComponents:1 vial of 100 μg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 μg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)Storage:The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at - 80°C.Preparation:HEK293T cells in 10-cm dishes were transiently transfected with MegaTran Transfection
Synonyms:DUP; FAP4; MRP1Predicted MW:127.2 kDaComponents:1 vial of 100 μg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 μg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)Storage:The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at - 80°C.Preparation:HEK293T cells in 10-cm dishes were transiently transfected with MegaTran Transfection
Predicted MW:127.2 kDaComponents:1 vial of 100 μg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 μg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)Storage:The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at - 80°C.Preparation:HEK293T cells in 10-cm dishes were transiently transfected with MegaTran Transfection
<ul> <li>Components: 1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)</li> <li>Storage: The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at - 80°C.</li> <li>Preparation: HEK293T cells in 10-cm dishes were transiently transfected with<u>MegaTran</u> Transfection</li> </ul>
<ul> <li>1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer</li> <li>1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)</li> <li>Storage: The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at -80°C.</li> <li>Preparation: HEK293T cells in 10-cm dishes were transiently transfected with<u>MegaTran</u> Transfection</li> </ul>
dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at - 80°C.Preparation:HEK293T cells in 10-cm dishes were transiently transfected with MegaTran Transfection
48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). Cell lysates were aliquoted and stored at -20°C before shipping.
<b>RefSeq:</b> <u>NP 002430</u>
Locus ID: 4437



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

SH3 (NM_002439) Human Over-expression Lysate – LY419328	
Cytogenetics:	5q14.1
Protein Families:	Druggable Genome, Stem cell - Pluripotency
Protein Pathways:	Colorectal cancer, Mismatch repair, Pathways in cancer
Product images:	



Western blot validation of overexpression lysate (Cat# LY419328) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with [RC212391] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US