

Product datasheet for LY419652

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

D Amino Acid Oxidase (DAO) (NM_001917) Human Over-expression Lysate

Product data:

Product Type: Over-expression Lysates

Description: Transient overexpression lysate of D-amino-acid oxidase (DAO)

Species: Human
Expression Host: HEK293T

Expression cDNA Clone

TrueORF Clone RC206768

or AA Sequence:

Tag: C-Myc/DDK

Detection Antibodies: Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)

ACCN: <u>NM 001917, NP 001908</u>

Synonyms: DAAO; DAMOX; OXDA

Predicted MW: 39.3 kDa

Components: 1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer

1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer

1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%

Bromophenol blue, 100mM DTT)

Storage: The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after

dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at -

80°C.

Preparation: HEK293T cells in 10-cm dishes were transiently transfected with MegaTran Transfection

Reagent (TT200002) and 5ug <u>TrueORF</u> cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). Cell lysates were aliquoted and stored at -20°C

before shipping.

RefSeq: NP 001908

Locus ID: 1610






Cytogenetics: 12q24.11

Protein Families: Druggable Genome

Protein Pathways: Arginine and proline metabolism, D-Arginine and D-ornithine metabolism, Glycine, serine and

threonine metabolism, Metabolic pathways

Product images:



Western blot validation of overexpression lysate (Cat# LY419652) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with [RC206768] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).