

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for LY423001

## FUNDC2P2 (NM\_001013648) Human Over-expression Lysate

## **Product data:**

| Product Type:                            | Over-expression Lysates  |  |
|--|--|--|
| Description:                             | Transient overexpression lysate of similar to hepatitis C virus core-binding protein 6; cervical cancer oncogene 3 (LOC388965)   |  |
| Species:                                 | Human  |  |
| Expression Host:                         | HEK293T  |  |
| Expression cDNA Clone<br>or AA Sequence: | TrueORF Clone RC209848   |  |
| Tag:                                     | C-Myc/DDK  |  |
| <b>Detection Antibodies:</b>             | Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)  |  |
| ACCN:                                    | <u>NM 001013648</u> , <u>NP 001013670</u>  |  |
| Synonyms:                                | MGC39809   |  |
| Predicted MW:                            | 16.9 kDa   |  |
| Components:                              | 1 vial of 100 μg gene specific transient over-expression cell lysate in RIPA buffer<br>1 vial of 100 μg whole HEK293T cell lysate in RIPA buffer<br>1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%<br>Bromophenol blue, 100mM DTT)  |  |
| Storage:                                 | The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at -80°C.   |  |
| Preparation:                             | HEK293T cells in 10-cm dishes were transiently transfected with <u>MegaTran</u> Transfection<br>Reagent (TT200002) and 5ug <u>TrueORF</u> cDNA plasmid. Transfected cells were cultured for<br>48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6,<br>150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF<br>and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was<br>measured by BCA kit (Thermo Scientific Inc.). Cell lysates were aliquoted and stored at -20°C<br>before shipping. |  |
| RefSeq:                                  | <u>NP_001013670</u>  |  |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

|               | FUNDC2P2 (NM_001013648) Human Over-expression Lysate – LY423001 |   |
|---------------|---|---|
| Locus ID:     | 388965  |   |
| Cytogenetics: | 2p11.2  |   |
| Product ima   | iges:   |   |
|               | 158-  |   |
|               | 106 —<br>79 —   | Western blot validation of overexpression lysate<br>(Cat# LY423001) using anti-DDK antibody (Cat#   |
|               | 48 -  | [TA50011-100]). Left: Cell lysates from un-<br>transfected HEK293T cells; Right: Cell lysates<br>from HEK293T cells transfected with [RC209848] |
|               | 35-<br>23-  | using transfection reagent MegaTran 2.0 (Cat#<br>[TT210002]).   |

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US