

## **Product datasheet for LY423465**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### STAR (NM 001007243) Human Over-expression Lysate

#### **Product data:**

**Product Type:** Over-expression Lysates

Description: Transient overexpression lysate of steroidogenic acute regulatory protein (STAR), nuclear

gene encoding mitochondrial protein, transcript variant 2

Species: Human **Expression Host:** HEK293T

**Expression cDNA Clone** 

or AA Sequence:

TrueORF Clone RC213299

C-Myc/DDK Tag:

**Detection Antibodies:** Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)

ACCN: NM 001007243, NP 001007244

Synonyms: cholesterol trafficker; mitochondrial steroid acute regulatory protein; StAR-related lipid

transfer (START) domain containing 1; STARD1; START domain containing 1; steroid acute

regulatory protein; steroidogenic acute regula; steroidogenic acute regulator

Predicted MW: 29.9 kDa

1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer Components:

1 vial of 100 μg whole HEK293T cell lysate in RIPA buffer

1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002%

Bromophenol blue, 100mM DTT)

Storage: The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after

> dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at -

80°C.

Preparation: HEK293T cells in 10-cm dishes were transiently transfected with MegaTran Transfection

> Reagent (TT200002) and 5ug TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). Cell lysates were aliquoted and stored at -20°C

before shipping.






**RefSeq:** NP 001007244

**Locus ID:** 6770

Cytogenetics: 8p11.23

**Protein Families:** Druggable Genome

# **Product images:**



Western blot validation of overexpression lysate (Cat# LY423465) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with [RC213299] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).