

Product datasheet for **MC200552**

Ndufa2 (NM_010885) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Ndufa2 (NM_010885) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Ndufa2
Synonyms:	AV000592; B8; C1-B8; CI-B8
Mammalian Cell Selection:	Neomycin
Vector:	PCMV6-Kan/Neo (PCMV6KN)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>BC006815 sequence for NM_010885 CAAAGATGGCGGCTGCCGCTGCTAGCCGAGCGGTCGGCGCAAAGCTGGGGTTGCGTGAGATTCGCGTTCA CTTATGCCAGCGTTCCCCAGGCAGCCAGGGTGTGAGGGATTTTCATCGTGCAACGGTACGTGGAGCTGAAG AAGGCGCACCCCAACCTGCCATTCTGATCCGCGAATGCTCGGAGGTGCAGCCCAAGCTTTGGGCCCGCT ATGCTTTTGGCCAAGAGAAGACGGTGTCTCTGAACAATCTGAGTGCTGATGAGGTAACCAGAGCCATGCA GAATGTGCTAAGCGGCAAGCCTGAAGGTCTCCACTGAGGACTGTGAGCGAGAGCAGCTGAACCTGCTGG ACTGAAGACAGTGTGGGAAATGTGTGCTTTGGGTCCTTATAAAGCTTACGCTGTACAGTAAAAAAAAA AAAAAA
Restriction Sites:	RsrII-NotI
ACCN:	NM_010885
Insert Size:	300 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



[View online »](#)

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [BC006815](#), [AAH06815](#)

RefSeq Size: 426 bp

RefSeq ORF: 300 bp

Locus ID: 17991

UniProt ID: [Q9CQ75](#)

Cytogenetics: 18 B2

Gene Summary: This gene encodes a subunit of the NADH-ubiquinone oxidoreductase (complex I) enzyme, which is a large, multimeric protein. It is the first enzyme complex in the mitochondrial electron transport chain and catalyzes the transfer of electrons from NADH to the electron acceptor ubiquinone. The proton gradient created by electron transfer drives the conversion of ADP to ATP. The human ortholog of this gene has been characterized, and its structure and redox potential is reported to be similar to that of thioredoxins. It may be involved in regulating complex I activity or assembly via assistance in redox processes. In humans, mutations in this gene are associated with Leigh syndrome, an early-onset progressive neurodegenerative disorder. A pseudogene of this gene is located on chromosome 5. [provided by RefSeq, May 2013]