

Product datasheet for MC200873

Chmp4b (BC011429) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Chmp4b (BC011429) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Chmp4b
Synonyms:	2010012F05Rik; C76846; Snf7-2
Mammalian Cell Selection:	Neomycin
Vector:	PCMV6-Kan/Neo (PCMV6KN)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>BC011429

```

CCCAGCTGCGGGGCGGAGGTGGAGGCGAGGCCTGCGGCTGCGGGGAGCGGCAGGCCGGGAGTGGGCGCGG
GAGCCGAGCCGAGCCGAGCCGAGCCGAGTGCGGCGCGGAGGCCGCGGCGAGCAGCAACCATGTCGG
TGTTTCGGGAAGCTGTTTCGGGGCTGGAGGGGGTAAGGCGGGCAAGGCGGCGCCGACCCCGAGGAGCCAT
CCAGCGGCTTCGGGACACGGAGGAGATGTTAAGCAAGAAGCAGGAGTTCTGGAGAAGAAAATCGAACAG
GAGCTGACGGCTGCCAAGAAGCACGGCACCAAAAATAAGCGCGCCGCCCTGCAGGCTCTGAAGCGCAAGA
AGAGGTATGAGAAGCAGCTGGCACAATTGATGGCACCCTGTCAACCATCGAGTTCAGCGGGAGGCCCT
AGAGAACGCCAACCAACACGGAGGTGCTCAAGAACATGGGCTATGCCGCCAAGGCCATGAAGGCTGCC
CACGACAACATGGACATTGATAAGGTGGATGAGTTAATGCAGGACATTGCTGACCAGCAAGAACTTGCAG
AGGAGATTTCCACAGCTATCTCCAAACCTGTGGGCTTTGGAGAAGAGTTCGACGAGGATGAGCTCATGGC
AGAGTTGGAGGAACCTGAACAAGAGGAGTTGGACAAGAATTTGTTGGAGATCAGTGGGCCCGAAACAGTC
CCTCTACCAAATGTCCCTCCGTAGCCCTACCATCCAAACCCGCCAAGAAGAAGGAAGAGGAAGATGACG
ACATGAAGGAATTGGAGAACTGGGCCGGATCCATGTAAGTGTTCAGCAGAGGCTGGGCCCGACAGCGACT
CTGGTGGCCTGTGCATCGGGCAGGCATGTGCGTGCGCAGGGCAGGCAGGACGCGGTGCAGGCAGCCTCCA
TCGCTCAGCTCTACCCAAAGCAGTAGCCGCACCACTGCTCACTCTCGCATAGCATGGTCTGTGCCCAGG
GGTGGGTGGGGGAGGGGGGCGGGGGGAGGTGCCTGCTGTTTATAATGTTGAATTTCTGTAATAATAAC
TGTATTTGCAAATCCAAAAAAAAAAAAAAAAA
  
```

Restriction Sites: RsrII-NotI

ACCN: BC011429

Insert Size: 675 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).


[View online »](#)

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	BC011429 , AAH11429
RefSeq Size:	1080 bp
RefSeq ORF:	675 bp
Locus ID:	75608
Cytogenetics:	2 H1
Gene Summary:	<p>Probable core component of the endosomal sorting required for transport complex III (ESCRT-III) which is involved in multivesicular bodies (MVBs) formation and sorting of endosomal cargo proteins into MVBs. MVBs contain intraluminal vesicles (ILVs) that are generated by invagination and scission from the limiting membrane of the endosome and mostly are delivered to lysosomes enabling degradation of membrane proteins, such as stimulated growth factor receptors, lysosomal enzymes and lipids. The MVB pathway appears to require the sequential function of ESCRT-O, -I, -II and -III complexes. ESCRT-III proteins mostly dissociate from the invaginating membrane before the ILV is released. The ESCRT machinery also functions in topologically equivalent membrane fission events, such as the terminal stages of cytokinesis. Together with SPAST, the ESCRT-III complex promotes nuclear envelope sealing and mitotic spindle disassembly during late anaphase. Plays a role in the endosomal sorting pathway. ESCRT-III proteins are believed to mediate the necessary vesicle extrusion and/or membrane fission activities, possibly in conjunction with the AAA ATPase VPS4. When overexpressed, membrane-assembled circular arrays of CHMP4B filaments can promote or stabilize negative curvature and outward budding. CHMP4A/B/C are required for the exosomal release of SDCBP, CD63 and syndecan.[UniProtKB/Swiss-Prot Function]</p>