

Product datasheet for MC200922

Psmb6 (NM_008946) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Psmb6 (NM_008946) Mouse Untagged Clone

Tag: Tag Free
Symbol: Psmb6

Synonyms: Lmp19; Mpnd

Mammalian Cell

Selection:

Neomycin

Vector: PCMV6-Kan/Neo (PCMV6KN)

E. coli Selection: Kanamycin (25 ug/mL)

Fully Sequenced ORF: >BC013897 sequence for NM_008946

Restriction Sites: Rsrll-Notl **ACCN:** NM 008946

Insert Size: 609 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>BC013897</u>, <u>AAH13897</u>

 RefSeq Size:
 805 bp

 RefSeq ORF:
 609 bp

 Locus ID:
 19175

 UniProt ID:
 Q60692

Cytogenetics: 11 42.99 cM

Gene Summary: Component of the 20S core proteasome complex involved in the proteolytic degradation of

most intracellular proteins. This complex plays numerous essential roles within the cell by associating with different regulatory particles. Associated with two 19S regulatory particles, forms the 26S proteasome and thus participates in the ATP-dependent degradation of ubiquitinated proteins. The 26S proteasome plays a key role in the maintenance of protein

homeostasis by removing misfolded or damaged proteins that could impair cellular

functions, and by removing proteins whose functions are no longer required. Associated with

the PA200 or PA28, the 20S proteasome mediates ubiquitin-independent protein degradation. This type of proteolysis is required in several pathways including

spermatogenesis (20S-PA200 complex) or generation of a subset of MHC class I-presented antigenic peptides (20S-PA28 complex). Within the 20S core complex, PSMB6 displays a peptidylglutamyl-hydrolyzing activity also termed postacidic or caspase-like activity, meaning that the peptides bond hydrolysis occurs directly after acidic residues. [UniProtKB/Swiss-Prot

Function]