

Product datasheet for **MC201434**

Isg15 (NM_015783) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Isg15 (NM_015783) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Isg15
Synonyms:	100038882; G1p2; IGI15; IP17; Irfp; UCRP
Mammalian Cell Selection:	Neomycin
Vector:	PCMV6-Kan/Neo (PCMV6KN)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>BC031424 sequence for NM_015783 CAGTCTGACTGTGAGAGCAAGCAGCCAGAAGCAGACTCCTTAATTCCAGGGGACCTAGAGCTAGAGCC TGCAGCAATGGCCTGGGACCTAAAGGTGAAGATGCTGGGGGTAACGATTCCTGGTGTCCGTGACTAAC TCCATGACGGTGTGAGAAGTGAAGAAGCAGATTGCCAGAAGATTGGTGTGCCGGCTTCCAGCAGCGCC TGGCCACCAAAGTGCAGTGCTCCAGGACGGTCTTACCCTTCCAGTCTGGGTCTGGGTCCAGCAGCAC AGTGATGCTAGTGGTACAGAAGTGCAGCGAGCCTCTGAGCATCCTGGTGGGAACGAAAGGGGCCACAGC AACATCTATGAGGTCTTCTGACGCAGACTGTAGACACGCTTAAGAAGAAGGTGTCCAGCGGGAACAAG TCCACGAAGACCAGTTCTGGCTGAGCTTCGAGGGAAGGCCCATGGAGGACAAGGAGCTGCTGGGGGAGTA TGGCCTAAAGCCCAGTGCACAGTGATCAAGCATTGCGCCTGAGGGGTGGGGGAGGGGACCAGTGTGCC TAAGGCCCCATCCAGAGCCAGTGTGGTTATAAAAGATGATGGGGGGGGGTATGGAGGACTTTTGGGA TAGCATTGAAAATGTAACGAGGAAAATACCTAATTAGAAAAAATATCAAAAAAAAAAAAAAAAAA
Restriction Sites:	RsrII-NotI
ACCN:	NM_015783
Insert Size:	486 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



[View online »](#)

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [BC031424](#), [AAH31424](#)

RefSeq Size: 694 bp

RefSeq ORF: 486 bp

Locus ID: 100038882

UniProt ID: [Q64339](#)

Cytogenetics: 4

Gene Summary: Ubiquitin-like protein which plays a key role in the innate immune response to viral infection either via its conjugation to a target protein (ISGylation) or via its action as a free or unconjugated protein. ISGylation involves a cascade of enzymatic reactions involving E1, E2, and E3 enzymes which catalyze the conjugation of ISG15 to a lysine residue in the target protein. Its target proteins include SERPINA3G/SPI2A, JAK1, MAPK3/ERK1, PLCG1, TRIM25, STAT5A, MAPK1/ERK2 and globin. Can also isgylate: DDX58/RIG-I which inhibits its function in antiviral signaling response and EIF4E2 which enhances its cap structure-binding activity and translation-inhibition activity. Exhibits antiviral activity towards both DNA and RNA viruses, including influenza A and B virus, sindbis virus (SV) and herpes simplex type-1 (HHV-1). Plays a significant role in the control of neonatal Chikungunya virus (CHIKV) infection by acting as a putative immunomodulator of proinflammatory cytokines. Protects mice against the consequences of Chikungunya virus infection by downregulating the pathogenic cytokine response, often denoted as the cytokine storm. Plays a role in erythroid differentiation. The secreted form of ISG15 can: induce natural killer cell proliferation, act as a chemotactic factor for neutrophils and act as a IFN-gamma-inducing cytokine playing an essential role in antimycobacterial immunity. The secreted form acts through the integrin ITGAL/ITGB2 receptor to initiate activation of SRC family tyrosine kinases including LYN, HCK and FGR which leads to secretion of IFNG and IL10; the interaction is mediated by ITGAL (By similarity). [UniProtKB/Swiss-Prot Function]