

## **Product datasheet for MC201815**

## Pla2g16 (NM\_139269) Mouse Untagged Clone

## **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** Pla2g16 (NM\_139269) Mouse Untagged Clone

Tag: Tag Free Symbol: Pla2g16

Synonyms: C78643; Hrasls3; Hrev107; HRSL3; MLP-3

Mammalian Cell

Selection:

Neomycin

Vector: PCMV6-Kan/Neo (PCMV6KN)

E. coli Selection: Kanamycin (25 ug/mL)

Fully Sequenced ORF: >BC024581 sequence for NM\_139269

**Restriction Sites:** RsrII-NotI **ACCN:** NM\_139269

**Insert Size:** 489 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Reconstitution Method:**

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** <u>BC024581</u>, <u>AAH24581</u>

RefSeq Size: 739 bp
RefSeq ORF: 489 bp
Locus ID: 225845
UniProt ID: Q8R3U1
Cytogenetics: 19 A

**Gene Summary:** Exhibits both phospholipase A1/2 and acyltransferase activities (PubMed:19047760). Shows

phospholipase A1 (PLA1) and A2 (PLA2), catalyzing the calcium-independent release of fatty

acids from the sn-1 or sn-2 position of glycerophospholipids (PubMed:18614531,

PubMed:19047760, PubMed:19136964, PubMed:22134920). For most substrates, PLA1 activity is much higher than PLA2 activity (By similarity). Shows O-acyltransferase activity, catalyzing the transfer of a fatty acyl group from glycerophospholipid to the hydroxyl group of lysophospholipid (By similarity). Shows N-acyltransferase activity,catalyzing the calcium-independent transfer of a fatty acyl group at the sn-1 position of phosphatidylcholine (PC) and other glycerophospholipids to the primary amine of phosphatidylethanolamine (PE), forming N-acylphosphatidylethanolamine (NAPE), which serves as precursor for N-

acylethanolamines (NAEs) (PubMed:19047760). Exhibits high N-acyltransferase activity and

low phospholipase A1/2 activity (By similarity).[UniProtKB/Swiss-Prot Function]