

Product datasheet for MC202028

Pqbp1 (NM_019478) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Pqbp1 (NM_019478) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Pqbp1
Synonyms:	npw38; PQBP-1; Sfc2
Mammalian Cell Selection:	Neomycin
Vector:	PCMV6-Kan/Neo (PCMV6KN)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>BC051673 sequence for NM_019478

```

AGAGCCTGTTGTGCTTTAAACCAAGAGCCCTTCGGTGTGGTCTGGTCAGCTACCCACTATGCCGCTTCC
TGTTGCGCTGCAGACCCGCTTGGCGAAGAGGAGCATCCTCAAACATCTGGAGCCGGAGCCAGAGGAAGAG
ATTATTGCTGAAGACTACGATGATGATCCTGTTGACTATGAGGCCACCCGATAGAGGGTCTGCCACCGA
GCTGGTACAAGGTGTTTGACCTTCTTGCAGACTCCCTTACTATTGGAATGTGGAGACAGACCTTGTGTC
GTGGCTCTCACCACATGATCCTAAGTTTGTGCTTACCAATCCGCAAGAAAGTCAGGAACAATAATGCA
GATGCTGAGGACAAGTCGGACCGGAATCTTGAAGAGGTGGACAGAAATCATGAGAAGTCAGATCGTAGTC
ATGAGAAGCCAGACAGGAGCCACGAGAAGGCAGACCGAAACCACGAGAAGAATGACAGAGAACGAGAGCG
CAACTACGACAAAGTGGATAGAGAGAGAGATCGGGACAGGGAACGAGAGCGGGCATTGACAAGGCAGAC
CGGAAGAGGGCAAGACCGACGTCACCATCGCAGAGAGGAAGTGGCTCCTTACCCCAAGAACAAGAAAG
CGACGAGCCGCAAGATGAAGAATTAGACCCCATGGACCCAGCTCATACTCAGATGCACCCCGGGGCAC
ATGGTCAACAGGACTCCCAAGAGGAACGAGGCCAAGACAGGTGCTGACACCACGGCAGCTGGGCCCTC
TTCCAGCAGCGCCCTTACCTTCCCCGGGCGCTGTGCTCCGCGCCAATGCAGAAGCTCCCGAACCAAAAC
AGCAGGACTGACTGACCCCTTGTTCCTGTAGCTTTGGCAGTTTGCCTTCTGTGTTTCTTCTTGTGGC
TCTGGGACAGGAGCCATGGCTTCTATGTGCTGCATTATTATCCACAGTGAGCTACACTCCCGGCCCT
TTAGTGTGTTTGTAAACAACAAGCCCTGTTGTTTAAATAAAAGCTTTTTTGTGGAACATAAAAAAAA
AAAAAAAAAAAAAAAAAAAA
  
```

Restriction Sites:	RsrII-NotI
ACCN:	NM_019478
Insert Size:	792 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).


[View online »](#)

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	BC051673 , AAH51673
RefSeq Size:	1070 bp
RefSeq ORF:	792 bp
Locus ID:	54633
UniProt ID:	Q91VJ5
Cytogenetics:	X 3.56 cM
Gene Summary:	<p>Intrinsically disordered protein that acts as a scaffold, and which is involved in different processes, such as pre-mRNA splicing, transcription regulation, innate immunity and neuron development (By similarity). Interacts with splicing-related factors via the intrinsically disordered region and regulates alternative splicing of target pre-mRNA species (PubMed:23512658). May suppress the ability of POU3F2 to transactivate the DRD1 gene in a POU3F2 dependent manner (By similarity). Can activate transcription directly or via association with the transcription machinery (By similarity). May be involved in ATXN1 mutant-induced cell death (By similarity). The interaction with ATXN1 mutant reduces levels of phosphorylated RNA polymerase II large subunit (By similarity). Involved in the assembly of cytoplasmic stress granule, possibly by participating to the transport of neuronal RNA granules (By similarity). Also acts as an innate immune sensor of infection by retroviruses, by detecting the presence of reverse-transcribed DNA in the cytosol (By similarity). Directly binds retroviral reverse-transcribed DNA in the cytosol and interacts with CGAS, leading to activate the cGAS-STING signaling pathway, triggering type-I interferon production (By similarity). [UniProtKB/Swiss-Prot Function]</p> <p>Transcript Variant: This variant (1) represents the longest transcript. Variants 1 and 2 encode the same isoform (1). Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.</p>