

Product datasheet for MC202082

Timp1 (NM 011593) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Timp1 (NM_011593) Mouse Untagged Clone

Tag: Tag Free
Symbol: Timp1

Synonyms: Clgi; EPA; Timp; TIMP-1; TPA-S1

Mammalian Cell

Selection:

Neomycin

Vector: PCMV6-Kan/Neo (PCMV6KN)

E. coli Selection: Kanamycin (25 ug/mL)

Fully Sequenced ORF: >BC051260 sequence for NM_011593

GAACACTGTCTACTTTTCCTCCATCTTTCTTTCTTTAGATGGTGAAATAAAGAACTATCAGACAGCAAA AAAAAAAAA

Restriction Sites: RsrII-NotI **ACCN:** NM_011593

Insert Size: 618 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>BC051260</u>, <u>AAH51260</u>

 RefSeq Size:
 780 bp

 RefSeq ORF:
 618 bp

 Locus ID:
 21857

 UniProt ID:
 P12032

 Cytogenetics:
 X 16.38 cM

Gene Summary: Metalloproteinase inhibitor that functions by forming one to one complexes with target

metalloproteinases, such as collagenases, and irreversibly inactivates them by binding to their catalytic zinc cofactor. Acts on MMP1, MMP2, MMP3, MMP7, MMP8, MMP9, MMP10, MMP11, MMP12, MMP13 and MMP16. Does not act on MMP14 (By similarity). Also functions as a growth factor that regulates cell differentiation, migration and cell death and activates

cellular signaling cascades via CD63 and ITGB1. Plays a role in integrin signaling.

[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (2) differs in the 5' UTR compared to variant 1. Variants 1 and 2

encode the same isoform (a).