

Product datasheet for MC202310

Eny2 (NM_175009) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Eny2 (NM_175009) Mouse Untagged Clone

Tag: Tag Free Symbol: Eny2

Synonyms: 1810057B09Rik; 6720481I12; DC6; Ey2

Mammalian Cell

Selection:

Neomycin

Vector: PCMV6-Kan/Neo (PCMV6KN)

E. coli Selection: Kanamycin (25 ug/mL)

Fully Sequenced ORF: >BC048361 sequence for NM_175009

AAAAAAAAAAAAAAA

Restriction Sites: Rsrll-Notl **ACCN:** NM_175009

Insert Size: 306 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>BC048361</u>, <u>AAH48361</u>

 RefSeq Size:
 647 bp

 RefSeq ORF:
 306 bp

 Locus ID:
 223527

 UniProt ID:
 Q9JIX0

 Cytogenetics:
 15 B3.2

Gene Summary: Involved in mRNA export coupled transcription activation by association with both the TREX-2

and the SAGA complexes. The transcription regulatory histone acetylation (HAT) complex SAGA is a multiprotein complex that activates transcription by remodeling chromatin and mediating histone acetylation and deubiquitination. Within the SAGA complex, participates in a subcomplex that specifically deubiquitinates both histones H2A and H2B. The SAGA complex is recruited to specific gene promoters by activators such as MYC, where it is required for transcription. Required for nuclear receptor-mediated transactivation. As a component of the TREX-2 complex, involved in the export of mRNAs to the cytoplasm through

the nuclear pores (By similarity).[UniProtKB/Swiss-Prot Function]