

## Product datasheet for MC202914

### Ppp1ca (NM\_031868) Mouse Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Ppp1ca (NM_031868) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Ppp1ca
Synonyms:	dism2; Ppp1c
Mammalian Cell Selection:	Neomycin
Vector:	PCMV6-Kan/Neo (PCMV6KN)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>BC014828 sequence for NM_031868

```

GGAGGCAGGAGAGGGCCCGAGCTGGTGGGCCGAGCGGCGGCCGCCATGTCCGACAGCGAGAAGCTC
AACCTGGACTCCATCATCGGGCGCCTGCTGGAAGTGCAGGGCTCACGGCCTGGGAAGAAGTGCAGCTGA
CAGAGAACGAGATCCGTGGTCTGTGCCTCAAATCCCGGGAGATTTCTCTGAGCCAGCCCATTTCTCTGGA
GCTTGAGGCGCCCTCAAGATCTGTGGTACATCCATGGCCAGTACTATGACCTTCTACGGCTGTTTGAG
TATGGTGGCTTCCCTCCAGAGAGCAACTACCTCTTCTGGGGGATTATGTAGATCGGGGCAAGCAGTCTT
TGGAGACCATCTGCCTGTTGCTGGCCTATAAGATCAGATACCCGGAAGATTTCTTCTACTTCTGTTGGAA
CCATGAGTGTGCCAGCATCAACCGCATTTATGGCTTCTATGATGAATGCAAGAGAAGATACAACATCAAA
CTGTGGAAGACGTTCACTGACTGCTTCACTGCCTGCCATTGCAGCCATTGTGGATGAGAAGATCTTCT
GCTGCCACGGGGCCTGTCTCCAGACTTGCAATCCATGGAGCAGATTAGGCGTATTATGCGGCCACAGA
CGTGCCTGACCAGGGCCTACTGTGTGATCTCCTGTGGTCTGACCCTGACAAGGATGTTCAAGGCTGGGGC
GAGAATGACCGTGGTGTCTCCTTTACCTTTGGGGCTGAGGTGGTAGCCAAGTTCTCTGCACAAGCATGATT
TGGACCTCATCTGCAGAGCACATCAGGTTGTAGAAGATGGCTATGAGTTCTTTGCCAAGAGACAGTTGGT
GACACTCTTCTCAGCTCCCAACTACTGTGGAGAGTTTGACAATGTGGTGCCATGATGAGTGTGGATGAG
ACCCTCATGTGTTCTTCCAGATCCTCAAGCCCGCTGATAAGAATAAGGGCAAGTATGGGCAAGTTCAGCG
GCCTGAACCCCGAGGGCCGCCATCACTCCACCCGCAATTCTGCCAAGCCAAGAAATAGCCTCCATG
TGCTGCCCTTCTGCCCAGATCGTTTGTACAGAAATCATGCTGCCATGGGTCACACTGGCCTCTCAGGCC
CACCCGTACGGGGAACACAGCGTTAAGTGTCTTTCTTTATTTTTTAAAGAAATCAATAGCAGCATCT
AATCTCCCAGGGCTCCCTCCCACAGCACCTGTGGTGGCTGCAAGTGAATCCTGGGGCCAAGGCTGCAG
CTCAGGGCAATGCGAGACCAGATTGTGGTCTCCAGCCTTGATGGCTGGCAGCCAGATCCTGGGGCAAC
CCATCTGGTCTCTTGAATAAAGGTCAAAGCTGGATTCTCAAAAAAAAAAAAAAAAAAAAAA
  
```

Restriction Sites:	RsrII-NotI
ACCN:	NM_031868
Insert Size:	993 bp


[View online »](#)

**OTI Disclaimer:** Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at [custsupport@origene.com](mailto:custsupport@origene.com) or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**Note:** Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.

**RefSeq:** [BC014828](#), [AAH14828](#)

**RefSeq Size:** 1392 bp

**RefSeq ORF:** 993 bp

**Locus ID:** 19045

**UniProt ID:** [P62137](#)

**Cytogenetics:** 19 A

**Gene Summary:**

Protein phosphatase that associates with over 200 regulatory proteins to form highly specific holoenzymes which dephosphorylate hundreds of biological targets. Protein phosphatase 1 (PP1) is essential for cell division, and participates in the regulation of glycogen metabolism, muscle contractility and protein synthesis. Involved in regulation of ionic conductances and long-term synaptic plasticity. May play an important role in dephosphorylating substrates such as the postsynaptic density-associated Ca(2+)/calmodulin dependent protein kinase II. Component of the PTW/PP1 phosphatase complex, which plays a role in the control of chromatin structure and cell cycle progression during the transition from mitosis into interphase. Regulates NEK2 function in terms of kinase activity and centrosome number and splitting, both in the presence and absence of radiation-induced DNA damage. Regulator of neural tube and optic fissure closure, and enteric neural crest cell (ENCCs) migration during development. In balance with CSNK1D and CSNK1E, determines the circadian period length, through the regulation of the speed and rhythmicity of PER1 and PER2 phosphorylation. May dephosphorylate CSNK1D and CSNK1E. Dephosphorylates CENPA (By similarity). Dephosphorylates the 'Ser-139' residue of ATG16L1 causing dissociation of ATG12-ATG5-ATG16L1 complex, thereby inhibiting autophagy (By similarity).[UniProtKB/Swiss-Prot Function]