

## Product datasheet for **MC202936**

### Zfp36 (NM\_011756) Mouse Untagged Clone

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** Zfp36 (NM\_011756) Mouse Untagged Clone  
**Tag:** Tag Free  
**Symbol:** Zfp36  
**Synonyms:** Gos24; Nup475; Tis11; TIS11D; TISII; Ttp; Zfp-36  
**Mammalian Cell Selection:** Neomycin  
**Vector:** PCMV6-Kan/Neo (PCMV6KN)  
**E. coli Selection:** Kanamycin (25 ug/mL)

**Fully Sequenced ORF:** >BC021391 sequence for NM\_011756  
 CCACGCGTCCGCAAGGCCATTTCGCGCCACCATGGATCTCTCTGCCATCTACGAGAGCCTCCAGTCGATGA  
 GCCATGACCTGTCATCCGACCACGGAGGAACCGAATCCCTCGGAGGACTTTGGAACATAAACTCGGACTC  
 CATCCCGTCTGGGGTACCTCTCGCCTGACTGGCCGCTCCACTAGCCTGGTGGAGGGCCGAAGCTGTGGC  
 TGGGTACCCCCACCCCTGGTTTTGCACCTTTGGCTCCCCGCCAGGCCCTGAGCTGTCACCCTACCTA  
 CTTTCGCTACTGCAACTCCCACCACCTCCTCTCGATACAAGACCGAGCTCTGTGGACCTACTCAGAAA  
 CGGGCGTTGTGCTACGGGGCAAGTGCCAGTTTGTCTACGGCCCTGGGTGAACTTCGCCAAGCCAATCGC  
 CACCCCAAGTACAAAACGGAAGTCTGCCACAAGTTCTACCTCCAGGGCCGCTGCCCTATGGCTCTCGAT  
 GCCACTTCATCCACAACCCACCGAGGACCTAGCTCTCCCTGGCCAGCCCATGTGCTGCGACAAAGCAT  
 CAGCTTCTCCGGTTGCCCTCAGGCCGAGAAGCTCGCCGCCACCTCCAGGCTTTTCTGGCCCTTCCCTG  
 TCCTCTTGTTCCTTTTCGCTTCCAGCTCCCCACCGCCCTGGGGATCTTCCACTTTCCCTTCTGCCT  
 TCTCTGCTGCCCTGGGACCCCTGTGACTCGAAGAGACCCTAACCCAGGCCTGTTGCCCTCTGCCGAAG  
 GTCTACTACCCCCAGCACCATCTGGGGGCCCTTGGGTGGCCTGGCTCGGAGCCCATCTGCCACTCTCTG  
 GGATCCGATCCTGATGACTACGCCAGCAGCGGCAGCAGCCTGGGGGGTTCAGACTCACCTGTCTTTGAGG  
 CAGGGGTGTTGGGCCTCCCCAGACCCCTGCACCCCAAGGCGTCTCCCATCTTCAATCGTATCTCTGT  
 CTCTGAGTGACAAGTGCCTACCTACCCAGTATGGATCAGCTAGATCTCAAAGAGAGGGCAGGGACTGCTC  
 ATTGCTGTGGGGACCTGGGGCACTCCTTAAGTTAATAAGTCCCCTCTTGGACATTCGAAGATGCAAT  
 AACCCATTTCCCTGGTGTGGCTGGGGCAGGTCCTAGTTTGCAAATTCAGTGTGGTGGATCCGTT  
 CCTAGGGTACCTAAGATGTTTGAGGGAGACAGTTGACAGTTGGTCTCCAGGCCCAAGTCTTCTGTTGT  
 TTTTGAGATAGGAGCTTATTATGGTACCCAGGCTGGCTTTGAACTCAATAAATCCTGCCTTAGCCTTT  
 TCCAAGTTCTGGGGTTACAGGTATGCACCAGCCCTCTGCAACTCTGGTCTCCTGGAATCTTAAGTGCTG  
 TGAAGAGCCGGCTCCCACAATACTATCCTAATTTTTACTAGACCCTGAAGTTCAGTGTCCGGTGGTCCGAA  
 GCCTCTCTGAGAATCCTGGTGTCAAATTTCCCTCCTAAAGCAAATAGCCAAAGCCATTGCCAAATCCC  
 TTCTCCCCAACCCAGTGGGCCCTTATTTATGACGACTTATTTATTGTATTAAGATTTTATAGTATTTA  
 TATATATTGGGTCGTCTACTCCGTTTTTCTTTTTGTAATGTTAAAAGTACTGATTAAGTATATGCT  
 ATAATATATTAATATATTGCTACAAAAAAAAAAAAAAAAAAAA



[View online »](#)

<b>Restriction Sites:</b>	RsrII-NotI
<b>ACCN:</b>	NM_011756
<b>Insert Size:</b>	960 bp
<b>OTI Disclaimer:</b>	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">BC021391</a> , <a href="#">AAH21391</a>
<b>RefSeq Size:</b>	1720 bp
<b>RefSeq ORF:</b>	960 bp
<b>Locus ID:</b>	22695
<b>UniProt ID:</b>	<a href="#">P22893</a>
<b>Cytogenetics:</b>	7 16.72 cM
<b>Gene Summary:</b>	Zinc-finger RNA-binding protein that destabilizes numerous cytoplasmic AU-rich element (ARE)-containing mRNA transcripts by promoting their poly(A) tail removal or deadenylation, and hence provide a mechanism for attenuating protein synthesis (PubMed:10330172, PubMed:10706852, PubMed:10805719, PubMed:15014438, PubMed:15187092, PubMed:15634918, PubMed:17030620, PubMed:19188452, PubMed:20595389, PubMed:21078877, PubMed:22701344, PubMed:27193233). Acts as an 3'-untranslated region (UTR) ARE mRNA-binding adapter protein to communicate signaling events to the mRNA decay machinery (PubMed:21278420). Recruits deadenylase CNOT7 (and probably the CCR4-NOT complex) via association with CNOT1, and hence promotes ARE-mediated mRNA deadenylation (PubMed:21278420). Functions also by recruiting components of the cytoplasmic RNA decay machinery to the bound ARE-containing mRNAs (PubMed:21278420). Self regulates by destabilizing its own mRNA (PubMed:15187092, PubMed:17288565). Binds to 3' UTR ARE of numerous mRNAs and of its own mRNA (PubMed:11533235, PubMed:15187092, PubMed:16508014, PubMed:17288565, PubMed:17971298, PubMed:20595389, PubMed:21078877, PubMed:21278420, PubMed:22701344, PubMed:27193233). Plays a role in anti-inflammatory responses; suppresses tumor necrosis factor (TNF)-alpha production by stimulating ARE-mediated TNF-alpha mRNA decay and

several other inflammatory ARE-containing mRNAs in interferon (IFN)- and/or lipopolysaccharide (LPS)-induced macrophages (PubMed:8630730, PubMed:9703499, PubMed:15014438, PubMed:16514065). Plays also a role in the regulation of dendritic cell maturation at the post-transcriptional level, and hence operates as part of a negative feedback loop to limit the inflammatory response (By similarity). Promotes ARE-mediated mRNA decay of hypoxia-inducible factor HIF1A mRNA during the response of endothelial cells to hypoxia (By similarity). Positively regulates early adipogenesis of preadipocytes by promoting ARE-mediated mRNA decay of immediate early genes (IEGs) (PubMed:22701344). Negatively regulates hematopoietic/erythroid cell differentiation by promoting ARE-mediated mRNA decay of the transcription factor STAT5B mRNA (By similarity). Plays a role in maintaining skeletal muscle satellite cell quiescence by promoting ARE-mediated mRNA decay of the myogenic determination factor MYOD1 mRNA (PubMed:25815583). Associates also with and regulates the expression of non-ARE-containing target mRNAs at the post-transcriptional level, such as MHC class I mRNAs (By similarity). Participates in association with argonaute RISC catalytic components in the ARE-mediated mRNA decay mechanism; assists microRNA (miRNA) targeting ARE-containing mRNAs (By similarity). May also play a role in the regulation of cytoplasmic mRNA decapping; enhances decapping of ARE-containing RNAs, in vitro (By similarity). Involved in the delivery of target ARE-mRNAs to processing bodies (PBs) (By similarity). In addition to its cytosolic mRNA-decay function, affects nuclear pre-mRNA processing (PubMed:22844456). Negatively regulates nuclear poly(A)-binding protein PABPN1-stimulated polyadenylation activity on ARE-containing pre-mRNA during LPS-stimulated macrophages (PubMed:22844456). Also involved in the regulation of stress granule (SG) and P-body (PB) formation and fusion (PubMed:15967811). Plays a role in the regulation of keratinocyte proliferation, differentiation and apoptosis (By similarity). Plays a role as a tumor suppressor by inhibiting cell proliferation in breast cancer cells (By similarity). [UniProtKB/Swiss-Prot Function]