

Product datasheet for MC202936

Zfp36 (NM_011756) Mouse Untagged Clone

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Expression Plasmids
Product Name:	Zfp36 (NM_011756) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Zfp36
Synonyms:	Gos24; Nup475; Tis11; TIS11D; TISII; Ttp; Zfp-36
Mammalian Cell Selection:	Neomycin
Vector:	PCMV6-Kan/Neo (PCMV6KN)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>BC021391 sequence for NM_011756 CCACGCGTCCGCAAGGCCATTCGCGCCACCATGGATCTCTCTGCCATCTACGAGAGCCTCCAGTCGATGA GCCATGACCTGTCATCCGACCACGGAGGAACCGAATCCCTCGGAGGACCTTTGGAACATAAACTCGGACTC CATCCCGTCTGGGGTCACCTCTCGCCTGACTGGCCGCTCCACTAGCCTGGGAGGGCCGAAGCTGTGGC TGGGTACCCCCACCCCCTGGTTTTGCACCTTTGGCTCCCCGCCCAGGCCTGAGCTGTCACCCTCACCTA CTTCGCCTACTGCAACTCCCACCACCTCCTCTCGATACAAGACCGAGCTCTGCCGAACCTACTCAGAAAG CGGGCGTTGTCGCTACGGGGCCAAGTGCCAGTTTGCTCACGGCCTGGGTGAACTTCGCCAAGCCAATCGC CACCCCAAGTACAAAACGGAACTCTGCCACAAGTTCTACCTCCAGGGCCGTGCCCCTATGGCTCTCGAT GCCACTTCATCCACAACCCCACCGAGGACCTAGCTCACCTGGCCAGCCCCAGGCCGCTGCCCCTATGGCTCTCGAT CACCTTCATCCCGGCTTGCCCTCAGGCCGCAGAAGCTCGCCCCCGGGGGATCTTCCACTTTCGGCCTTCCTG TCCTCTGTTCCTTTTCGCCTTCCAGGCCGCCAAGAGCCCCCGGGGGATCTTCCACTTTCCGCCTTCTGCT CCTCTGGCCCCCAGGACCCCTGGGCCCTGGGTGGCCTGGCTCGGAGCCCACTCGCCAGCACAAGCAT CAGGTTTCGGCCCTCCCAGACCGCGCCCTGGGGGGGCCCGGCCCACTCGCCCCCCTGCGGAG GTCTACTACCCCCAGCACCATCTGGGGGCCCTGGCGGGGGGCCCGACTCGCCCCCTCTGG GGATCCGATCC



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE Zfp36 (NM_011756) Mouse Untagged Clone – MC202936

Restriction Sites:	Rsrll-Notl
ACCN:	NM_011756
Insert Size:	960 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	 Centrifuge at 5,000xg for 5min. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	<u>BC021391, AAH21391</u>
RefSeq Size:	1720 bp
RefSeq ORF:	960 bp
Locus ID:	22695
UniProt ID:	<u>P22893</u>
Cytogenetics:	7 16.72 cM

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Zfp36 (NM_011756) Mouse Untagged Clone - MC202936

Gene Summary:

Zinc-finger RNA-binding protein that destabilizes numerous cytoplasmic AU-rich element (ARE)-containing mRNA transcripts by promoting their poly(A) tail removal or deadenylation, and hence provide a mechanism for attenuating protein synthesis (PubMed:10330172, PubMed:10706852, PubMed:10805719, PubMed:15014438, PubMed:15187092, PubMed:15634918, PubMed:17030620, PubMed:19188452, PubMed:20595389, PubMed:21078877, PubMed:22701344, PubMed:27193233). Acts as an 3'-untranslated region (UTR) ARE mRNA-binding adapter protein to communicate signaling events to the mRNA decay machinery (PubMed:21278420). Recruits deadenylase CNOT7 (and probably the CCR4-NOT complex) via association with CNOT1, and hence promotes ARE-mediated mRNA deadenylation (PubMed:21278420). Functions also by recruiting components of the cytoplasmic RNA decay machinery to the bound ARE-containing mRNAs (PubMed:21278420). Self regulates by destabilizing its own mRNA (PubMed:15187092, PubMed:17288565). Binds to 3' UTR ARE of numerous mRNAs and of its own mRNA (PubMed:11533235, PubMed:15187092, PubMed:16508014, PubMed:17288565, PubMed:17971298, PubMed:20595389, PubMed:21078877, PubMed:21278420, PubMed:22701344, PubMed:27193233). Plays a role in anti-inflammatory responses; suppresses tumor necrosis factor (TNF)-alpha production by stimulating ARE-mediated TNF-alpha mRNA decay and several other inflammatory ARE-containing mRNAs in interferon (IFN)- and/or lipopolysaccharide (LPS)-induced macrophages (PubMed:8630730, PubMed:9703499, PubMed:15014438, PubMed:16514065). Plays also a role in the regulation of dendritic cell maturation at the post-transcriptional level, and hence operates as part of a negative feedback loop to limit the inflammatory response (By similarity). Promotes ARE-mediated mRNA decay of hypoxia-inducible factor HIF1A mRNA during the response of endothelial cells to hypoxia (By similarity). Positively regulates early adipogenesis of preadipocytes by promoting ARE-mediated mRNA decay of immediate early genes (IEGs) (PubMed:22701344). Negatively regulates hematopoietic/erythroid cell differentiation by promoting ARE-mediated mRNA decay of the transcription factor STAT5B mRNA (By similarity). Plays a role in maintaining skeletal muscle satellite cell quiescence by promoting ARE-mediated mRNA decay of the myogenic determination factor MYOD1 mRNA (PubMed:25815583). Associates also with and regulates the expression of non-ARE-containing target mRNAs at the posttranscriptional level, such as MHC class I mRNAs (By similarity). Participates in association with argonaute RISC catalytic components in the ARE-mediated mRNA decay mechanism; assists microRNA (miRNA) targeting ARE-containing mRNAs (By similarity). May also play a role in the regulation of cytoplasmic mRNA decapping; enhances decapping of ARE-containing RNAs, in vitro (By similarity). Involved in the delivery of target ARE-mRNAs to processing bodies (PBs) (By similarity). In addition to its cytosolic mRNA-decay function, affects nuclear pre-mRNA processing (PubMed:22844456). Negatively regulates nuclear poly(A)-binding protein PABPN1-stimulated polyadenylation activity on ARE-containing pre-mRNA during LPSstimulated macrophages (PubMed:22844456). Also involved in the regulation of stress granule (SG) and P-body (PB) formation and fusion (PubMed:15967811). Plays a role in the regulation of keratinocyte proliferation, differentiation and apoptosis (By similarity). Plays a role as a tumor suppressor by inhibiting cell proliferation in breast cancer cells (By similarity). [UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US