

Product datasheet for **MC203258**

Siah1a (NM_009172) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Siah1a (NM_009172) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Siah1a
Synonyms:	AA982064; AI853500; Sinh1a
Mammalian Cell Selection:	Neomycin
Vector:	PCMV6-Kan/Neo (PCMV6KN)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>BC046317

```

CCGGGCTGCCGGGCGGGCGCGCTCTCGAGGGCGGCGGCCAGGGTGTCCCGTCGGTCTCGGTGCCGGGAG
AGGTGGAGGCGCGGCCCGCGGTGGCGCGGTGACGGTGGCGGTGGCGGGCGGCGGACGTGGCGA
CCGGGCGCGTTAGACGGGGCGGGGTCCGCGGCGCGCTCTCCGCCACAGAGATGAGCCGCCAGACTGCT
ACAGCATTACCCACTGGCACCTCAAAGTGTCCACCATCCAGAGGGTACCTGCCTTGACCGGCACAACTG
CATCCAACAATGACTTGGCGAGTCTTTTGTAGTGTCTGTCTGCTTTGACTATGTGTTGCCACCTATTCT
TCAGTGTGAGAGTGGCCATCTTGTGTGTAGCAACTGTGCGCCCAAACTTACATGTTGTCCCACTTGCCGG
GGCCCATTTGGGATCCATTGCAACTTGGCTATGGAGAAAGTGGCCAACTCAGTACTCTTCCCTTGTAAT
ATGCTCTTCTGGATGTGAAATAACTCTGCCACACACCGAAAAGGCAGAGCACGAGGAGCTCTGTGAGTT
CAGGCCTTACTCCTGCCCTGCCCTGGTCTTCTGTAAAGTGGCAAGGCTCCTTGGATGCCGTGATGCC
CACCTGATGCATCAGCACAAGTCCATTACCACCCTGCAAGGAGAAGATATAGTTTTCCTTGCTACAGACA
TTAACCTTCTGTGTGCTGTTGACTGGGTGATGATGCACTTGTGTTTGGCTTTCATTTTCATGTTAGTCTT
GGAGAAACAAGAAAAATATGATGGTCATCAGCAGTTCTTTGCAATTGTACAACATGATAGGAACACGCAAG
CAAGCTGAAAAATTTGCTATCGACTTGAGCTAAATGGTCATAGGCGGCGATTGACTTGGGAAGCGACTC
CTCGGTCTATTGATGAGGGAATTGCAACAGCCATTATGAATAGTACTGCCTAGTGTGTTGACACCAAGCAT
TGCACAGCTTTTGCAGAAAAATGGCAATTTAGGCATCAATGTAATATTTCCATGTGTTGAAACGGCAAT
CAAATATTTCTGGCCAGTGTGTTAAATTTGCATTTGACTTCACAGAGAATAAGGCACCCATCTGCTTGCC
AACCTAAAACTTCTCGGTAGGTAGAAGCTAGACATGAAGTAAATAAAAAAGAAAGCTGTTAATACAGG
AAACAGTTGCATGTAGTAACACTAATATATTTAAAAATAATTCAACAGTAAACCACTGAAAAAATATAT
ATACCCAAGATGGGCATCTTTTGTATTAAGAAAGGAAACATTGTAAAAATTTTCTGAACTTTGTGTTGT
TGTAGATTGATTGATTGTTGACAATTTTTTGGGGGTGTGTGTCTGTGCACGCATGCGTGACGTGTGTG
GTTGGTTTTCTTTAACTGACAAGCCATCTGCGTGGTCATAGACCACTGTTTTCCCTTGTGAGTCAACA
CATAGTGTGCTGTGGGTTTGTGTTTTCTGTTTTGTTTTGTTTTGATGTGTGTTTGTCTAA
TTTTTATTCTAGTTTTTCATTAATAAATTTGACTTCTTTCTGTAAAAAAGAAAAAAGAAAAA
AAAAAAGAAAAAAGAAAAAAGAAAAAAGAAAAAAGAAAAAAGAAAAAAGAAAAAAGAAAAAAGAAAAA

```

Restriction Sites: RsrII-NotI


[View online »](#)

ACCN:	NM_009172
Insert Size:	849 bp
OTI Disclaimer:	<p>Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.</p> <p>The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info</p>
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	BC046317 , AAH46317
RefSeq Size:	1645 bp
RefSeq ORF:	849 bp
Locus ID:	20437
UniProt ID:	P61092
Cytogenetics:	8 42.1 cM

Gene Summary:

E3 ubiquitin-protein ligase that mediates ubiquitination and subsequent proteasomal degradation of target proteins. E3 ubiquitin ligases accept ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfers the ubiquitin to targeted substrates. Mediates E3 ubiquitin ligase activity either through direct binding to substrates or by functioning as the essential RING domain subunit of larger E3 complexes. Triggers the ubiquitin-mediated degradation of many substrates, including proteins involved in transcription regulation (ELL2, MYB, POU2AF1, PML and RBBP8), a cell surface receptor (DCC), the cell-surface receptor-type tyrosine kinase FLT3, the cytoplasmic signal transduction molecules (KLF10/TIEG1 and NUMB), an antiapoptotic protein (BAG1), a microtubule motor protein (KIF22), a protein involved in synaptic vesicle function in neurons (SYP), a structural protein (CTNNB1) and SNCAIP. Confers constitutive instability to HIPK2 through proteasomal degradation. It is thereby involved in many cellular processes such as apoptosis, tumor suppression, cell cycle, axon guidance, transcription, spermatogenesis and TNF-alpha signaling. Has some overlapping function with SIAH2. Required for completion of meiosis I in males. Induces apoptosis in cooperation with PEG3. Upon nitric oxid (NO) generation that follows apoptotic stimulation, interacts with S-nitrosylated GAPDH, mediating the translocation of GAPDH to the nucleus. GAPDH acts as a stabilizer of SIAH1, facilitating the degradation of nuclear proteins.[UniProtKB/Swiss-Prot Function]