

## Product datasheet for MC203796

### Selenos (NM\_024439) Mouse Untagged Clone

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** Selenos (NM\_024439) Mouse Untagged Clone  
**Symbol:** Selenos  
**Synonyms:** 1500011E07Rik; C78786; H-4; H-47; H4; H47; Se; Sels; Seps1; V; Vimp  
**Mammalian Cell Selection:** Neomycin  
**Vector:** PCMV6-Kan/Neo (PCMV6KN)  
**E. coli Selection:** Kanamycin (25 ug/mL)  
**Fully Sequenced ORF:** >BC011091  
 CGCAGAGCAGAGAGGAGCAGCGGGCTTGCTGTTGGTGGCGGTGGCCGCCATGGATCGCGATGAG  
 GAACCTCTGTCTCCGAGGCCGGCGCTGGAGACCGAGAGCCTGCGATTCTGCACGTGACAGTGGGCTCCC  
 TGCTGGCCAGCTATGGCTGGTACATCCTCTTCAGCTGCATCCTACTCTACATTGTCATCCAGAGGCTCTC  
 CCTTCGACTGAGGGCTTTGAGGCAGAGACAGCTGGACCAAGCCGAGACTGTTCTGGAACCTGATGTTGTT  
 GTTAAGCGACAAGAGGCTTTAGCAGCTGCTCGTTTGAGAATGCAGGAAGATCTAAATGCCAAGTTGAAA  
 AACATAAGGAAAACTAAGACAGCTTGAAGAAGAAAAAGAACAGAGAAGATTGAAATGTGGGACAGCAT  
 GCAAGAAGGCAGAAGTTACAAAAGAAATTCAGGAAGGCCCTCAGGAAGAAGATGGTCTGGACCTTCTACT  
 TCATCTGTCATCCCCAAGGAAAATCTGACAAAAAGCCTTTGCGAGGAGGTGGTTATAACCCTCTGACGG  
 GTGAAGGGGGTGAACCTGCTCCTGGAGACCTGGACGCAGGGGCCCATCATCTGGTGGATGAACTAAGA  
 CTCTTGTTAGTGTGCTCTGACATTAGCAAGGTGAACCTTTAACCTCACTCAATTGCCTTACGCACAC  
 TTTCACAGTGACTGGCCAAGGAGAGGTGGGGCTTTTCTCTGTTCTAACTACTTGTACTTTAAGGGCTTT  
 GGTGAGCATGAGATATAGACATTGCCATTAGGCCACACTCTAGACAAGACAGCCATGGCTTTTCATGGCTG  
 CTGGCTAGTTGGTAGGTTGAAGGCTTCTTGCTGTTTAGCAGACTTCATAAAGGAGGCCAGTGATGATAC  
 TTTGGGTAGAAAGTCCTTGCTGACAGGATGGTCTCTGTGACGGGATGCGTTGAATGATGTCTTCTTATA  
 AATGGTGAACCCACAGTGAGGATTACTGATGTTACAGTTGATGGGGTTTGCTTCTGTATATTTATTTT  
 ATGTACAGAACTTTGTAAAAAAGTTAAATACTTAAAAAGTAACATTTTATGATCTTTATTAAC  
 TCAAGGAAATTTCTTTGTGAGCTTGACTTTGTCAGACAGTAAACAGCTTTGTATCAAAAAAAAAAAAAA  
 AAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

**Restriction Sites:** RsrII-NotI

**ACCN:** NM\_024439

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP). The expression of this clone is not guaranteed due to the nature of selenoproteins.


[View online »](#)

<b>OTI Annotation:</b>	This clone encodes a selenoprotein containing the rare amino acid selenocysteine (Sec). Sec is encoded by UGA codon, which normally signals translational termination. Expression of this clone is not guaranteed due to the nature of selenoproteins.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"> <li>1. Centrifuge at 5,000xg for 5min.</li> <li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>3. Close the tube and incubate for 10 minutes at room temperature.</li> <li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
<b>Note:</b>	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
<b>RefSeq:</b>	<u><a href="#">BC011091</a></u> , <u><a href="#">AAH11091</a></u>
<b>RefSeq Size:</b>	1219 bp
<b>Locus ID:</b>	109815
<b>UniProt ID:</b>	<u><a href="#">Q9BCZ4</a></u>
<b>Cytogenetics:</b>	7 35.49 cM
<b>Gene Summary:</b>	<p>This gene encodes a transmembrane protein that is localized in the endoplasmic reticulum (ER). It is involved in the degradation process of misfolded proteins in the ER, and may also have a role in inflammation control. This protein is a selenoprotein, containing the rare amino acid selenocysteine (Sec). Sec is encoded by the UGA codon, which normally signals translation termination. The 3' UTRs of selenoprotein mRNAs contain a conserved stem-loop structure, designated the Sec insertion sequence (SECIS) element, that is necessary for the recognition of UGA as a Sec codon, rather than as a stop signal. Two additional phylogenetically conserved stem-loop structures (Stem-loop 1 and Stem-loop 2) in the 3' UTR of this mRNA have been shown to function as modulators of Sec insertion (PMID:23614019). Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jul 2017]</p> <p>Transcript Variant: This variant (1) represents the predominant transcript and encodes the longer isoform (1).</p>