

Product datasheet for MC203864

Exd2 (NM_133798) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Exd2 (NM_133798) Mouse Untagged Clone

Tag: Tag Free Symbol: Exd2

Synonyms: 4930539P14Rik; C85658; Exdl2

Vector: PCMV6-Kan/Neo

E. coli Selection: Kanamycin (25 ug/mL)

Cell Selection:NeomycinRestriction Sites:Rsrll-NotlACCN:NM_133798

Insert Size: 1491 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: BC018508, AAH18508

RefSeq Size: 2865 bp
RefSeq ORF: 1491 bp
Locus ID: 97827



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Exd2 (NM_133798) Mouse Untagged Clone - MC203864

UniProt ID: Q8VEG4

Cytogenetics: 12 C3

Gene Summary: Exonuclease required for double-strand breaks resection and efficient homologous

recombination. Plays a key role in controlling the initial steps of chromosomal break repair, it is recruited to chromatin in a damage-dependent manner and functionally interacts with the MRN complex to accelerate resection through its 3'-5' exonuclease activity, which efficiently processes double-stranded DNA substrates containing nicks.[UniProtKB/Swiss-Prot Function]